

ON THE GEOMETRY OF TORIC ARRANGEMENTS

C. DE CONCINI AND C. PROCESI

Dedicated to Vladimir Drinfeld on the occasion of his 50th birthday

ABSTRACT. Motivated by the counting formulas of integral polytopes as in Brion–Vergne [5], [4] and Szenes–Vergne [26], we start to lay the foundations of a theory for toric arrangements, which may be considered as the *periodic version* of the theory of hyperplane arrangements.

1. INTRODUCTION

1.1. The knapsack problem. One of the aims of this paper is to understand in a more algebraic geometric language, the counting formulas of [5], [4], [26] for integral points in polytopes.

We are given a family Δ of vectors in a lattice Λ and, for $\beta \in \Lambda$ we want to count the number c_β of solutions of the equation:

$$\sum_{\alpha \in \Delta} n_\alpha \alpha = \beta, \quad n_\alpha \in \mathbb{N}.$$

This is clearly the value of a *partition function*, or in other words, the number of integral points of the convex polytope

$$\Pi_\beta := \{(x_\alpha) \mid \sum_{\alpha \in \Delta} x_\alpha \alpha = \beta, \quad 0 \leq x_\alpha \in \mathbb{R}\}.$$

One usually restricts to the case in which Δ is all on one side of a hyperplane so that the polytope is bounded and we have finitely many solutions. In this case the numbers c_β are finite and are the coefficients of the generating series:

$$(1) \quad f_\Delta := \sum_{\beta} c_\beta e^\beta = \prod_{\alpha \in \Delta} \frac{1}{1 - e^\alpha}$$

Thus the problem is to understand an *inversion formula*, the formula which computes these numbers c_β from f_Δ .

A prototype example is the *knapsack problem*, i.e. the same problem where now the elements α are just positive integers h_i , (cf. the paper of E.T. Bell [3]). In this setting, we ask, given an integer n to compute the coefficient of x^n in the function $\prod_i \frac{1}{1-x^{h_i}}$ or to compute the residue $\frac{1}{2\pi i} \oint \prod_i \frac{x^{-n-1}}{1-x^{h_i}} dx$ over a small circle. The function $\prod_i \frac{x^{-n-1}}{1-x^{h_i}}$ has poles in 0 and in the m -th

The authors are partially supported by the Cofin 40 %, MIUR.

roots of 1, where m is the least common multiple of the numbers h_i . In other words, if $\zeta = e^{2\pi i/m}$ one can write $\prod_i \frac{1}{1-x^{h_i}} = \prod_{i=1}^m \frac{1}{(1-\zeta^i x)^{b_i}}$ where the integers b_i are easily computed from the h_j . By standard residue theory we have:

$$(2) \quad \frac{1}{2\pi i} \oint \prod_i \frac{x^{-n-1}}{1-x^{h_i}} dx = - \sum_{j=1}^m \frac{1}{2\pi i} \oint_{C_j} \prod_{i=1}^m \frac{x^{-n-1}}{(1-\zeta^i x)^{b_i}} dx$$

where C_j is a small circle around ζ^{-j} .

We are using the fact that the residue at ∞ is 0, due to the hypothesis made that $n > 0$.

For the term $\frac{1}{2\pi i} \oint_{C_j} \prod_{i=1}^m \frac{x^{-n-1}}{(1-\zeta^i x)^{b_i}} dx$ one makes a coordinate change $x = w + \zeta^{-j}$ and obtains

$$\frac{1}{2\pi i} \oint_{C_j} \prod_{i=1}^m \frac{(w + \zeta^{-j})^{-n-1}}{(1 - \zeta^{i-j} - \zeta^i w)^{b_i}} dw.$$

Now $\prod_{i=1, i \neq j}^m \frac{1}{(1 - \zeta^{i-j} - \zeta^i w)^{b_i}}$ is holomorphic around 0 and so it can be expanded as a power series $\sum_{h=0}^{\infty} a_{j,h} w^h$ whose terms can be computed, while

$$(w + \zeta^{-j})^{-n-1} = \zeta^{j(n+1)} \sum_{k=0}^{\infty} (-1)^k \binom{n+k}{k} (\zeta^j w)^k.$$

Finally we have for the j -th term of (2):

$$\begin{aligned} -\frac{(-1)^{b_j}}{2\pi i} \oint_{C_j} \zeta^{j(n+1-b_j)} \left(\sum_{k=0}^{\infty} (-1)^k \binom{n+k}{k} \zeta^{jk} w^k \right) \left(\sum_{h=0}^{\infty} a_{j,h} w^h \right) w^{-b_j} dw = \\ -(-1)^{b_j} \zeta^{j(n+1-b_j)} \sum_{k+h=b_j-1} (-1)^k \zeta^{jk} \binom{n+k}{k} a_{j,h}. \end{aligned}$$

This formula, summed over all j , answers the question and exhibits the partition function requested as a sum of functions which are polynomials on the cosets modulo m . One calls such a function a *periodic polynomial* or *quasipolynomial*.

In this paper we show that the counting formulas of [5], [4], [26], which are the higher dimensional case of the knapsack problem, can be interpreted in a similar way as sums of residues at the *poles* of the generating function. In order to develop this higher dimensional case, we shall study a trigonometric analog of the theory of hyperplane arrangements, which we call a *toric arrangement*, in which the main object of study is the open set of a torus complement of a finite set of cosets of codimension 1 tori. Versions of these arrangements also known as toral arrangements have been introduced and studied in [11], [12], [17], [18], [19], [20], [21], in particular in [19], Looijenga proves, by a completely different argument, using the degeneration of a spectral sequence, that the cohomology of this space has a natural filtration whose graded spaces coincide with the ones we describe in Theorem 4.2,

in particular the Betti numbers are computed. As for the algebra structure, contrary to what claimed in [19], 2.4.3, the algebra is generated by the logarithmic differentials if and only if the set of characters is *unimodular*, see §5.2 and (5.2). Toric arrangements are the main topic of this paper.

There is a more analytic approach to counting formulas, via splines. It is due to Dahmen, Micchelli cf. [7], and it would be interesting to compare the two approaches.

Finally there is some overlap, at least in the development of expansion formulas, with the work of A. Szenes, [25].

1.2. Toric arrangements. Let $T = \text{hom}_{\mathbb{Z}}(\Lambda, \mathbb{C}^*)$ be an algebraic torus over $\mathbb{C}$, of dimension r . Λ its character group, a rank r lattice, $A := \mathbb{C}[T] = \mathbb{C}[\Lambda]$ its coordinate ring.

We consider, as analogue of a hyperplane, a translate of the kernel of a character χ , i.e. the hypersurface $H_{a,\chi}$ of equation $1 - a\chi = 0$ for a suitable non zero $a \in \mathbb{C}^*$.

Given a finite subset $\Delta \subset \mathbb{C}^* \times \Lambda$, we shall study the **components of the arrangement** $\mathcal{R}_{\Delta}$ generated by the hypersurfaces $H_{a,\chi}$ as $(a, \chi) \in \Delta$. By this we shall mean *the set of all connected components* of all intersections of these hypersurfaces.

The complement of the union of these hypersurfaces will be denoted by $\mathcal{A}_{\Delta}$, it is an affine variety of coordinate ring $A[d^{-1}]$, $d := \prod_{(a,\chi) \in \Delta} (1 - a\chi)$.

Our first concern will be to compute the cohomology of $\mathcal{A}_{\Delta}$. For this, let us remark that on the entire torus T we have the closed forms $d \log \chi$, $\chi \in \Lambda$, spanning the vector space of invariant 1-forms isomorphic to $\Lambda \otimes \mathbb{R}$ and generating the cohomology of T . On the open set $\mathcal{A}_{\Delta}$ we have then also the closed one forms $d \log(1 - a\chi)$, $(a, \chi) \in \Delta$. In general, unless the arrangement is *unimodular* (see 5.2), these two sets of 1-forms are not enough to generate cohomology (this corrects a statement in [19] 2.4.3).

Nevertheless one can give a satisfactory description, in term of explicit differential forms of $H^*(\mathcal{A}_{\Delta}, \mathbb{C})$. One obtains then:

Theorem 4.2. *For each integer $i \geq 0$, we have a (non canonical) decomposition, as W runs over the components of the arrangement:*

$$H^i(\mathcal{A}_{\Delta}) = \oplus_W H^{i-\text{codim}W}(W) \otimes V_W.$$

Each W is isomorphic to a torus, so its cohomology is an exterior algebra on r explicit forms, and V_W is a finite dimensional vector space, which depends on the combinatorics of the set Δ via the theory on non broken circuits and can be identified to the top cohomology of the hyperplane arrangement defined by the differentials, at a given point of W , of the functions $1 - a\chi$ vanishing on W .

The result is particularly striking in the unimodular case, where we shall prove *formality*, i.e.

Theorem 5.2 *The subalgebra of the algebra of differential forms generated by the 1-forms $d \log(1-a\chi)$, $d \log \chi$ is isomorphic to the De Rham cohomology of $\mathcal{A}_\Delta$.*

The algebraic relations between these forms resemble, but are more complicated, the relations of Orlik-Solomon (cf. 5.1) for the hyperplane case. The theory of non broken circuits furnishes a linear basis.

The proof of these results will be based on a careful analysis of $\mathcal{A}_\Delta$. We shall use some simple ideas from D -modules to do a precise bookkeeping.

The reader unfamiliar with D -modules can find, in the introductory book [6] of S. C. Coutinho, essentially all that we need.

We have limited the analysis to De Rham cohomology. It is likely that a finer geometric analysis will produce a similar result (after normalizing the classes), for integral cohomology.

1.3. Residues. The second task is to define correctly the residues which will be integrals over suitable cycles. Since we are in dimension > 1 we are faced with the problem that the poles are on divisors with a complicated intersection pattern, this implies that we need to use a model where the divisor has normal crossings as in [8]. This is done in §6.

Finally we have to understand which cycles contribute to the counting formula. This is done in §7, it is just a reinterpretation of results of Szenes–Vergne [26] and it is connected to the combinatorial problem of understanding the regions (chambers) over which the partition function is a periodic polynomial. The answer is expressed by the notion of the Jeffrey–Kirwan residue.

With this analysis the counting formulas are explained in a very similar way to the 1-dimensional case.

2. PERIODIC WEYL ALGEBRA

2.1. Differential operators. Given a torus T with character group Λ , we shall now consider the ring D_T of differential operators over T . D_T is generated by $A = \mathbb{C}[T] = \mathbb{C}[\Lambda]$ and by the invariant derivations D_ϕ , $\phi \in \Lambda^*$ which are defined by $D_\phi(\chi) := \langle \phi, \chi \rangle \chi$, for any $\chi \in \Lambda$. In terms of a basis of characters $\{\xi_1, \dots, \xi_m\}$, we have the basis of invariant derivations $\partial_i := \xi_i \frac{\partial}{\partial \xi_i}$.

We shall think of D_T as a periodic version of the Weyl algebra. In explicit coordinates the *periodic* Weyl algebra is

$$W^p(r) := \mathbb{C}[\xi_1^{\pm 1}, \dots, \xi_r^{\pm 1}, \frac{\partial}{\partial \xi_1}, \dots, \frac{\partial}{\partial \xi_r}] = \mathbb{C}[\xi_1^{\pm 1}, \dots, \xi_r^{\pm 1}, \partial_1, \dots, \partial_r]$$

If Λ is the character group of T we shall also denote $D_T = W(\Lambda)$.

We filter the algebra D_T by writing its elements as

$$u := \sum a_{i_1, \dots, i_r} \partial_1^{i_1} \dots \partial_r^{i_r}, \quad a_{i_1, \dots, i_r} \in A$$

and putting such an element in degree $\max \sum_{j=1}^r i_j$ for which the coefficient $a_{i_1, \dots, i_r}$ is non 0. The associated graded algebra is a polynomial ring $A[\eta_1, \dots, \eta_r]$ which can be viewed as the coordinate ring of the cotangent bundle to the torus T (trivialized by the invariant forms).

When we have a finitely generated module M one can define its *characteristic variety*, which is an important geometric invariant of the module, applying to D_T the same method used for modules over the Weyl algebra (cf. [6]). We define a *Bernstein filtration*, by choosing a finite set of generators m_i and setting $M_j := \sum_i (D_T)_j m_i$. The graded module associated is a module over $A[\eta_1, \dots, \eta_r]$.

The radical of its annihilator ideal is independent of the choice of the generators m_i so it defines a subvariety $ch(M)$ of the cotangent bundle of T the *characteristic variety of M* .

A basic fact on D -modules is that this variety is of dimension $\geq \dim M$ and, when it is of minimal dimension $= \dim M$, the module is said to be *holonomic*. It is again a basic fact, see again [6], that the coordinate ring of $\mathcal{A}_\Delta$ is a holonomic D -module and has a finite composition series. The task of the next section is to introduce and describe the irreducible modules which appear in this composition series as will finally be shown in Proposition 3.4. The special irreducible modules which appear can be thought of as the natural *quantization* of the cotangent bundles of the components of the arrangement. They have a very special description which allows us to put them together to produce explicit partial fraction decompositions for the elements of the coordinate ring of the affine variety $\mathcal{A}_\Delta$ i.e. of the localized ring $\mathbb{C}[T][d^{-1}]$, $d := \prod_{(a, \chi) \in \Delta} (1 - a\chi)$.

2.2. Special Modules. Take a sublattice $\Gamma \subset \Lambda$. Fix a maximal ideal $\mathfrak{m} \subset \mathbb{C}[\Gamma]$ and denote by $I_{\mathfrak{m}}$ the extension of $\mathfrak{m}$ to A . The variety $W \subset T$ defined by $I_{\mathfrak{m}}$ is a coset of the subgroup T_Γ kernel of the characters in Γ and will be called for short a *coset*.

Γ is recovered from W as the set of characters which are constant on W , clearly $\mathfrak{m}$ is formed by the elements of $\mathbb{C}[\Gamma]$ vanishing on W .

Set $\Gamma^\perp = \{\phi \in \Lambda^* | \phi(\Gamma) = 0\}$. Next let $J_{\mathfrak{m}}$ be the left ideal in D_T generated by $\mathfrak{m}$ and by the derivations D_ϕ with $\phi \in \Gamma^\perp$.

Remark The derivations D_ϕ with $\phi \in \Gamma^\perp$ are the invariant vector fields tangent to W .

We want to study the structure of the D_T module:

$$(3) \quad N(W) = N(\Gamma, J_{\mathfrak{m}}) := D_T / J_{\mathfrak{m}}.$$

We first treat the case in which Γ is a split direct summand, which is equivalent to saying that T_Γ and W are connected.

Under this assumption, we can then take a basis $\xi_1, \dots, \xi_r$ of the character group so that Γ is generated by $\xi_1, \dots, \xi_k$ for some k .

The derivations $\partial_i := \xi_i \frac{\partial}{\partial \xi_i}$ are a basis of the invariant vector fields. The ∂_i , $i = k+1, \dots, r$ are a basis of the space of the derivations D_ϕ with $\phi \in \Gamma^\perp$.

The ideal $\mathfrak{m} = (1 - a_1\xi_1, \dots, 1 - a_k\xi_k)$ for suitable $a_1, \dots, a_k \in \mathbb{C}^*$.

Lemma 2.1. *With the previous notations, and with W a connected coset:*

i) $N(W)$ is an irreducible D_T -module having as linear basis the classes of the elements

$$t_{\underline{h}, \underline{n}} := \xi_{k+1}^{h_{k+1}} \dots \xi_r^{h_r} \prod_{j=1, \dots, k} \frac{\partial^{n_j}}{\partial \xi_j^{n_j}}$$

$h_s \in \mathbb{Z}, n_j \in \mathbb{N}$.

Or alternatively the elements

$$u_{\underline{h}, \underline{n}} := \xi_{k+1}^{h_{k+1}} \dots \xi_r^{h_r} \prod_{j=1, \dots, k} \partial_j^{n_j}$$

$h_s \in \mathbb{Z}, n_j \in \mathbb{N}$.

ii) $N(W)$ is a holonomic module with characteristic variety the conormal bundle to W .

i) The elements $\prod_{s=k+1}^r \xi_s^{h_s} \prod_{j=1}^k \frac{\partial^{n_j}}{\partial \xi_j^{n_j}} \prod_{j=k+1}^r \frac{\partial^{n_j}}{\partial \xi_j^{n_j}} \prod_{s=1}^k \xi_s^{h_s}$ with $h_s \in \mathbb{Z}$ and $n_j \in \mathbb{N}$ constitute a basis for the ring D_T . This clearly implies that the classes of the $t_{\underline{n}, \underline{h}}$'s with $h_s \in \mathbb{Z}, n_j \in \mathbb{N}$, span $N(W)$.

It remains to show that the classes of the $t_{\underline{n}, \underline{h}}$'s with $h_s \in \mathbb{Z}, n_j \in \mathbb{N}$ are linearly independent and that $N(W)$ is irreducible. Denote by v the class of 1 modulo $J_{\mathfrak{m}}$. Take an element

$$w = \sum_{(\underline{h}, \underline{n}) \in S} a_{\underline{h}, \underline{n}} t_{\underline{h}, \underline{n}} v$$

where S is a subset of $\mathbb{Z}^{r-k} \times \mathbb{N}^k$ and the $a_{\underline{h}, \underline{n}}$ are non zero complex numbers. Multiplying by a suitable monomial in $\xi_{k+1}, \dots, \xi_r$, we can clearly assume that $S \subset \mathbb{N}^{r-k} \times \mathbb{N}^k = \mathbb{N}^r$. We want to show that the submodule M generated by w contains v . This will clearly implies both claims. We make induction on the largest element s of S in the lexicographic ordering. If this element is zero there is nothing to prove since w is a non zero multiple of v .

Otherwise, write $s = (s_1, \dots, s_r)$ and let $1 \leq i \leq r$ be the least element such that s_i is non zero. If $i \leq k$, then we get

$$(1 - a_i \xi_i) w = \sum_{(\underline{n}, \underline{h}) \in S} a_{\underline{h}, \underline{n}} [(1 - a_i \xi_i), t_{\underline{h}, \underline{n}}] v.$$

Using the fact that $[(1 - a_i \xi_i), \frac{\partial}{\partial \xi_i}^m] = m a_i \frac{\partial}{\partial \xi_i}^{m-1}$, we deduce that

$$(1 - a_i \xi_i) w = \sum_{(\underline{n}, \underline{h}) \in S'} a_{\underline{h}, \underline{n}} t_{\underline{h}, \underline{n}} v$$

with the maximum element in S' lexicographically smaller than the maximum element in S . Thus by induction we get that v lies in the D_T submodule generated by w .

If $i > k$, we apply $\frac{\partial}{\partial \xi_i}$ and, reasoning in exactly the same way, we deduce that also in this case M contains v .

For the other elements we use the fact that $\partial_i = \frac{\partial}{\partial \xi_i} \xi_i - 1$ and see that we have a triangular base change.

ii) When we compute the characteristic variety we see that the graded module associated to $N(W)$ is still cyclic so it equals $A[\eta_1, \dots, \eta_r]/J$ where J is the ideal generated by the elements $1 - a_j \xi_j$, $j = 1, \dots, k$, η_h , $h = k+1, \dots, r$ which is the conormal bundle to W . $\square$

Let us go back to a general $\Gamma \subset \Lambda$ and set $\bar{\Gamma} = \{\chi \in \Lambda \mid \exists n \text{ with } n\chi \in \Gamma\}$. $\bar{\Gamma}$ is a split direct summand in Λ . Let $h = |\bar{\Gamma}/\Gamma|$. The variety W associated to the ideal $I_{\mathfrak{m}}$ is now a union $W = \cup_{i=1}^h W_i$ of h connected components, cosets of the connected torus $T_{\bar{\Gamma}}$. These components are defined by the maximal ideals $\mathfrak{m}_1, \dots, \mathfrak{m}_h \subset \mathbb{C}[\bar{\Gamma}]$ lying over $\mathfrak{m}$. We can clearly order the set $\{\phi_1, \dots, \phi_h\}$ of primitive idempotents in $\mathbb{C}[\bar{\Gamma}]/\mathfrak{m}\mathbb{C}[\bar{\Gamma}]$ in such a way that the annihilator of ϕ_j in $\mathbb{C}[\bar{\Gamma}]$ is $\mathfrak{m}_j$ for each $j = 1, \dots, h$.

Remark $\bar{\Gamma}$ is the set of characters in Λ which are *locally constant* on W .

Lemma 2.2. *There is a canonical isomorphism of D_T modules,*

$$N(W) \simeq \oplus_{i=1}^h N(W_i).$$

Proof. Since clearly $J_{\mathfrak{m}} \subset J_{\mathfrak{m}_i}$ for each $i = 1, \dots, h$, we have a surjective homomorphism

$$f_i : N(W) \rightarrow N(W_i).$$

Taking the direct sum of the f_i 's, we then get a homomorphism

$$f : N(W) \rightarrow \oplus_{i=1}^h N(W_i).$$

Notice that since the $N(W_i)$'s are pairwise non isomorphic (having different characteristic varieties), we get that f is surjective.

On the other hand, if we denote by v the class of 1 in $N(W)$ and set $v_i = \phi_i v$ (notice that this makes sense since $\mathfrak{m}$ annihilates v), we have that $J_{\mathfrak{m}_i}$ annihilates v_i . We thus get homomorphisms

$$g_i : N(W_i) \rightarrow N(W)$$

$i = 1, \dots, h$. Taking their sum we get

$$g : \oplus_{i=1}^h N(W_i) \rightarrow N(W)$$

Since $v = \sum_{i=1}^h v_i$, we immediately deduce that g is surjective.

We leave to the reader the simple verification that f and g are one inverse to the other. $\square$

We are now going to give explicit realizations of the module $N(W)$.

Before doing this, we need a general statement which is well known but whose proof we include for completeness. Let R be a commutative ring. $\underline{z} = (z_1, \dots, z_k)$ a sequence of elements in R such that any permutation of them is a regular sequence. For any subset $S \subset \{1, \dots, k\}$, set $z_S = \prod_{j \notin S} z_j$. Consider the Čech complex $C(R, \underline{z})$

$$0 \rightarrow R \rightarrow \oplus_{i=1}^k R[z_i^{-1}] \rightarrow \dots \rightarrow \oplus_{S, |S|=t} R[z_S^{-1}] \rightarrow \dots \rightarrow R[(z_1 \cdots z_k)^{-1}] \rightarrow 0$$

Lemma 2.3. 1) $H^i(C(R, \underline{z})) = 0$ for $i = 0, \dots, k-1$.

2) The R submodule of $H^k(C(R, \underline{z}))$, generated by the class of $(z_1 \cdots z_k)^{-1}$ is isomorphic to $R/(z_1, \dots, z_k)$. In particular $H^k(C(R, \underline{z})) \neq 0$.

Proof. We proceed by induction on k . If $k = 1$ our complex is

$$0 \rightarrow R \rightarrow R[z^{-1}] \rightarrow 0$$

the map $R \rightarrow R[z^{-1}]$ is injective since z is not a zero divisor. It is not surjective since z is not invertible. Clearly the annihilator of the class of $1/z$ in $R[z^{-1}]/R$ consists of those elements $r \in R$ such that $r/z \in R$. This is the ideal (z) .

We now assume our statement for $k-1$. Consider the ring $R' = R/(z_k)$ and in it the sequence $\underline{z}' = (z'_1, \dots, z'_{k-1})$ where z'_j is the image of z_j modulo z_k . Our assumptions are satisfied by R' and the sequence $\underline{z}'$. Also we have an exact sequence of complexes

$$0 \rightarrow C(R, \underline{z}) \xrightarrow{\cdot z_k} C(R, \underline{z}) \rightarrow C(R', \underline{z}') \rightarrow 0.$$

Applying the inductive assumption and the long cohomology sequence we immediately deduce that $H^i(C(R, \underline{z})) = 0$ for $0 \leq i \leq k-2$ and multiplication by z_k gives an injective map

$$0 \rightarrow H^{k-1}(C(R, \underline{z})) \xrightarrow{\cdot z_k} H^{k-1}(C(R, \underline{z})).$$

We claim that z_k acts locally nilpotently on $H^{k-1}(C(R, \underline{z}))$ so that necessarily $H^{k-1}(C(R, \underline{z})) = 0$. Indeed let $(r_1, \dots, r_k) \in \oplus_{i=1}^k R[z_{\{i\}}]$ be a cycle. This means that $r_1 + r_2 + \dots + r_k = 0$. If we multiply by a big enough power of z_k , say z_k^M , we get that $z_k^M r_i \in R[z_{\{i,k\}}]$ for each $i = 1, \dots, k-1$. Then $z_k^M(r_1, \dots, r_k) = (z_k^M r_1, \dots, z_k^M r_k)$ is cohomologous to $(0, \dots, 0, z_k^M(r_1 + r_2 + \dots + r_k)) = (0, \dots, 0)$ as desired.

It remains to show 2). For this again using the long cohomology sequence and 1) we get an exact sequence

$$0 \rightarrow H^{k-1}(C(R', \underline{z}')) \xrightarrow{\delta} H^k(C(R, \underline{z})) \xrightarrow{\cdot z_k} H^k(C(R, \underline{z})) \rightarrow 0.$$

It is easy to see that $\delta((z'_1 \cdots z'_{k-1})^{-1}) = (z_1 \cdots z_k)^{-1}$. This and our inductive assumption immediately imply our claim (notice that $R/(z_1, \dots, z_k) = R'/(z'_1, \dots, z'_{k-1})$). $\square$

Remark 2.4. It is well known that $H^i(C(R, \underline{z}))$ equals the i -th local cohomology group $H^i_{(z_1, \dots, z_k)}(R)$ (see for example [14] where our Lemma 2.3 is essentially contained).

We are now ready to give explicit realizations of the module $N(W) = N(\Gamma; J_{\mathfrak{m}})$. To achieve this let us choose a basis $\Psi := \{\psi_1, \dots, \psi_k\}$ in Λ of Γ so that the ideal $\mathfrak{m} = (1 - a_1 \psi_1, \dots, 1 - a_k \psi_k)$ for suitable non zero constants $a_i \in \mathbb{C}^*$ defines W . Set $d = \prod_{i=1}^k (1 - a_i \psi_i)$. and for any $i = 1, \dots, k$, $d_i = d/(1 - a_i \psi_i)$. Consider the ring $A[d^{-1}]$ as D_T module and remark that $A[d_i^{-1}]$ is a submodule for each i . We have

Proposition 2.5. *The D_T module $A[d^{-1}]/\sum_{i=1}^k A[d_i^{-1}]$ is isomorphic to $N(W)$.*

Proof. As before let $\mathfrak{m}_1, \dots, \mathfrak{m}_h \subset \mathbb{C}[\bar{\Gamma}]$ be set set of maximal ideals lying over $\mathfrak{m}$, the set $\{\phi_1, \dots, \phi_h\}$ the set of primitive idempotents in $\mathbb{C}[\bar{\Gamma}]/m\mathbb{C}[\bar{\Gamma}]$ ordered in such a way that the annihilator of ϕ_j in $\mathbb{C}[\bar{\Gamma}]$ is $\mathfrak{m}_j$ for each $j = 1, \dots, h$.

Denote by w the class of d^{-1} in $A[d^{-1}]/\sum_{i=1}^k A[d_i^{-1}]$. It is immediate to see that $J_{\mathfrak{m}}$ annihilates w so that we get a map of D_T modules

$$f : N(\Gamma; J_{\mathfrak{m}}) \rightarrow A[d^{-1}]/\sum_{i=1}^k A[d_i^{-1}]$$

with $f(v) = w$. To see that this map is injective it suffices to show that its restriction to each irreducible component $N(\bar{\Gamma}; J_{\mathfrak{m}_i})$ is non zero. $N(\bar{\Gamma}; J_{\mathfrak{m}_i})$ is generated by $v_i = \phi_i v$. We have $f(v_i) = \phi_i f(v) = \phi_i w$ and this is non zero by lemma 2.3.

Thus we only have to show that f is surjective. $A[d^{-1}]/\sum_{i=1}^k A[d_i^{-1}]$ is spanned by the classes of the elements

$$\frac{a}{(1 - a_1 \psi_1)^{n_1} \dots (1 - a_k \psi_k)^{n_k}}$$

with $a \in A$ and $n_i > 0$ for each $i = 1, \dots, k$. In $\Lambda^* \otimes \mathbb{Q}$ choose elements $\rho_1, \dots, \rho_k$ such that $\langle \rho_j, \psi_i \rangle = \delta_{i,j}$ for each $i, j = 1, \dots, k$. Take the derivations $D_i := (a_i \psi_i)^{-1} D_{\rho_i}$. It is then immediate to see that

$$\frac{a D_1^{n_1} \dots D_k^{n_k}}{n_1! \dots n_k!} \left(\frac{1}{(1 - a_1 \psi_1) \dots (1 - a_k \psi_k)} \right) = \frac{a}{(1 - a_1 \psi_1)^{n_1} \dots (1 - a_k \psi_k)^{n_k}}$$

and everything follows. $\square$

3. TORIC ARRANGEMENTS

3.1. Toric arrangements. Let $T = \text{hom}_{\mathbb{Z}}(\Lambda, \mathbb{C}^*)$ be an algebraic torus over $\mathbb{C}$, of dimension r . Λ its character group, a rank r lattice, $A := \mathbb{C}[T] = \mathbb{C}[\Lambda]$ its coordinate ring.

We want to study in this setting a suitable analogue of the theory of hyperplane arrangements. We develop the theory in a level of generality higher than the one needed for the applications to counting formulas.

We consider, as analogue of a hyperplane, a translate of the kernel of a character χ , i.e. the hypersurface $H_{a,\chi}$ of equation $1 - a\chi = 0$ for a suitable non zero $a \in \mathbb{C}^*$.

Notice that, if χ is not primitive, this hypersurface is not connected, but it is the union of finitely many cosets of the codimension 1 torus, the connected component of the identity of the kernel of χ . If $\Psi := \{\chi_1, \dots, \chi_i\}$ are linearly independent characters and $\underline{a} = (a_1, \dots, a_i)$ is a sequence of non zero complex numbers, the intersection $K_{\underline{a}, \Psi}$ of the H_{a_s, χ_s} is in general not connected even if the characters are primitive.

In fact let us denote by $\langle \Psi \rangle := \langle \chi_1, \dots, \chi_i \rangle$ the subgroup generated by the $\chi_j, j = 1, \dots, i$, and by $\overline{\langle \Psi \rangle}$ the subgroup of Λ formed by the elements χ such that $\chi^m \in \langle \Psi \rangle$ for some $m > 0$. The following is immediate:

Proposition 3.1. *$K_{\underline{a}, \Psi}$ is a coset of the subgroup kernel of the characters in $\langle \Psi \rangle$. It has $|\overline{\langle \Psi \rangle} / \langle \Psi \rangle|$ connected components each of which is a coset of the subgroup kernel of the characters in $\overline{\langle \Psi \rangle}$.*

Given a finite subset $\Delta \subset \mathbb{C}^* \times \Lambda$, we shall study the components of the **arrangement** $\mathcal{R}_\Delta$ generated by the hypersurfaces $H_{a, \chi}$ as $(a, \chi) \in \Delta$.

As explained in the introduction, by this we shall mean the *set of all connected components* of all intersections of these hypersurfaces.

The complement of the union of these hypersurfaces will be denoted by $\mathcal{A}_\Delta$, it is an affine variety of coordinate ring $A[d^{-1}]$, $d := \prod_{(a, \chi) \in \Delta} (1 - a\chi)$. Our main concern will be to compute the cohomology of $\mathcal{A}_\Delta$.

3.2. The D_T -module R . With the notations of the previous section, if W is one component of the arrangement $\mathcal{R}_\Delta$, let $\Delta(W)$ be the set of elements $(a, \chi) \in \Delta$ with $1 - a\chi = 0$ on W and $\Sigma(W)$ the set of characters $\chi \in \Lambda$ which, as functions on W , are constant. It is easily seen that $\Sigma(W)$ is a split direct summand on Λ and that W is a coset of the subtorus $T(W)$ kernel of the characters of $\Sigma(W)$.

The subtori $T(W)$ will be called the *tori associated to the arrangement*.

The complement of the union of the hypersurfaces $H_{a, \chi}$ will be denoted by $\mathcal{A}_\Delta$, it is an affine variety of coordinate ring $R = A[d^{-1}]$, with $d := \prod_{(a, \chi) \in \Delta} (1 - a\chi)$. We begin by studying the ring R as a D_T module.

Let $\pi : \Delta \rightarrow \Lambda$ be the projection on the second factor and set $\overline{\Delta} := \pi(\Delta)$.

We shall say that a subset $S = \{(a_1, \psi_1), \dots, (a_r, \psi_r)\} \subset \Delta$ is linearly independent if $\{\psi_1, \dots, \psi_r\} \subset \Lambda$ are linearly independent. We set for each $k = 0, \dots, r$,

$$\mathcal{J}_k = \{S \subset \Delta \mid S \text{ is linearly independent and } |S| \leq k\}.$$

Of course $\mathcal{J}_{r+s} = \mathcal{J}_r := \mathcal{J}$ for each $s \geq 0$ (J_0 is by definition the empty set). Given $S \in \mathcal{J}$ we set $d_S := \prod_{(a, \chi) \in S} (1 - a\chi)$ ($d_\emptyset = 1$), and define for each $k = 0, \dots, r$,

$$R_k = \sum_{S \in \mathcal{J}_k} A[d_S^{-1}].$$

Clearly $R_k \subset R_{k+1}$ for each k and R_k is a sub D_T module of R .

The next Proposition is a special case of Proposition 4.6 of [25].

Proposition 3.2. *Let $\Psi \subset \mathbb{C}^* \times \Lambda$ be finite. Let $\Xi \subset \Lambda$ denote the projection of Ψ to the second factor and set Γ equal to the lattice generated by Ξ . Then*

$$\prod_{(a, \psi) \in \Psi} \frac{1}{(1 - a\psi)}$$

can be written as a linear combination with coefficients in the ring of $\mathbb{C}[\Gamma]$ of elements of the form

$$\frac{1}{(1 - a_1\psi_1)^{h_1} \cdots (1 - a_r\psi_r)^{h_r}}$$

where $(a_i, \psi_i) \in \Psi$ for each $i = 1, \dots, r$ and $\{\psi_1, \dots, \psi_r\}$ are linearly independent.

Proof. By a simple induction we may assume that $\Psi = \{(a_0, \psi_0), \dots, (a_n, \psi_n)\}$ with $\psi_0, \dots, \psi_n$ linearly dependent and $\psi_1, \dots, \psi_n$ linearly independent. We then choose representatives $\phi_i \in \mathbb{C}[\Gamma]$ of the primitive idempotents e_i of the ring $\mathbb{C}[\Gamma]/(1 - a_1\psi_1, \dots, 1 - a_n\psi_n)$. Notice that in this ring we have $\psi_0 e_i = \beta_i e_i$ with $\beta_i \in \mathbb{C}$. By the definition of the ϕ_i 's, we have $1 = \sum_i \phi_i + \sum_{j=1}^n b_i(1 - a_j\psi_j)$, for some $b_i \in \mathbb{C}[\Gamma]$. So

$$\begin{aligned} \frac{1}{(1 - a_0\psi_0) \cdots (1 - a_n\psi_n)} &= \frac{\sum_i \phi_i + \sum_{j=1}^n b_i(1 - a_j\psi_j)}{(1 - a_0\psi_0) \cdots (1 - a_n\psi_n)} = \\ &= \sum_i \frac{\phi_i}{(1 - a_0\psi_0) \cdots (1 - a_n\psi_n)} + \sum_{j=1}^n \frac{b_i(1 - a_j\psi_j)}{(1 - a_0\psi_0) \cdots (1 - a_n\psi_n)} \end{aligned}$$

Using induction on the cardinality of Ψ we then have to analyze only the terms

$$\frac{\phi_i}{(1 - a_0\psi_0) \cdots (1 - a_n\psi_n)}.$$

We have that $(1 - a_0\psi_0)\phi_i = \sum_{j=1}^n c_j(1 - a_j\psi_j) + \gamma_i\phi_i$ with $\gamma_i = 1 - a_0\beta_i$. We separate two cases. If $\gamma_i = 0$, we have $(1 - a_0\psi_0)\phi_i = \sum_{j=1}^n c_j(1 - a_j\psi_j)$ and substituting

$$\frac{\phi_i}{(1 - a_0\psi_0) \cdots (1 - a_n\psi_n)} = \frac{\sum_{j=1}^n c_j(1 - a_j\psi_j)}{(1 - a_0\psi_0)^2(1 - a_1\psi_1) \cdots (1 - a_n\psi_n)}$$

we obtain a sum of terms in which in the denominator some $1 - a_j\psi_j$ has disappeared.

If $\gamma_i \neq 0$, we get

$$\frac{\phi_i}{(1 - a_0\psi_0) \cdots (1 - a_n\psi_n)} = \gamma_i^{-1} \frac{(1 - a_0\psi_0)\phi_i - \sum_{j=1}^n c_j(1 - a_j\psi_j)}{(1 - a_0\psi_0) \cdots (1 - a_n\psi_n)},$$

and again everything follows by induction. $\square$

3.3. Partial fractions for R . From Proposition 3.2 we deduce

Corollary 3.3. $R = R_r$.

Proof. The proof is clear. $\square$

Let us now take $S \subset \mathcal{J}_k$. Let $\Gamma \subset \Lambda$ be the sublattice generated by the characters $\pi(S)$, $\mathfrak{m}$ the maximal ideal in $\mathbb{C}[\Gamma]$ generated by the elements $1 - a\chi$ with $(a, \chi) \in S$. As in the previous section introduce the lattice $\overline{\Gamma} = \{\chi \in \Lambda \mid \exists n \text{ with } \chi^n \in \Gamma\}$. Set $h = |\overline{\Gamma}/\Gamma|$. Let $\mathfrak{m}_1, \dots, \mathfrak{m}_h \subset \mathbb{C}[\overline{\Gamma}]$ be the set of maximal ideals lying over $\mathfrak{m}$, and order the set $\{\phi_1, \dots, \phi_h\}$ of primitive

idempotents in $\mathbb{C}[\overline{\Gamma}]/m\mathbb{C}[\overline{\Gamma}]$ in such a way that the annihilator of ϕ_j in $\mathbb{C}[\overline{\Gamma}]$ is $\mathfrak{m}_j$ for each $j = 1, \dots, h$.

From Proposition 2.5 we deduce a canonical isomorphism as D_T -modules:

$$(4) \quad A[d_S^{-1}]/(R_{k-1} \cap A[d_S^{-1}]) \simeq N(\Gamma; J_{\mathfrak{m}}) \simeq \bigoplus_{i=1}^h N(\overline{\Gamma}; J_{\mathfrak{m}_i})$$

mapping the class of 1 in $N(\overline{\Gamma}; J_{\mathfrak{m}_i})$ to $\phi_i[1/d_S]$, with $[1/d_S]$ the class of the vector $1/d_S$.

Let W be a component of the arrangement of codimension k . We shall say that $S \subset \mathcal{J}_k$ and W are associated if W is a component of the variety defined by the vanishing of the elements $1 - a\chi$ with $(a, \chi) \in S$. In this case denoting by $\mathfrak{n}_S$ the ideal generated by the elements $1 - a\chi$ with $(a, \chi) \in S$ we have that $\mathbb{C}[\Lambda]/\mathfrak{n}_S$ has a primitive idempotent $\phi_{W,S}$ which, as function on Z_S is the characteristic function of the component W .

We associate to such a W a subspace $V_W \subset R_k/R_{k-1}$ as follows.

Definition 1. Define V_W as the subspace spanned by the classes $v_{S,W} := \phi_{W,S}[1/d_S]$ as S runs among the subsets of $\mathcal{J}_k$ associated to W .

Notice that the $\mathbb{C}[\Lambda]$ module generated by d_S^{-1} in R_k/R_{k-1} factors as an $\mathbb{C}[\Lambda]/\mathfrak{n}_S$ module.

Proposition 3.4. For each $k = 0, \dots, r$, R_k/R_{k-1} is a semisimple D_T module and we have a canonical isomorphism

$$(5) \quad R_k/R_{k-1} \simeq \bigoplus_W N(W) \otimes V_W$$

W runs over the components of the arrangement of codimension k .

Proof. R_k/R_{k-1} is a quotient of $\bigoplus_{S \in \mathcal{J}_k} A[d_S^{-1}]/(R_{k-1} \cap A[d_S^{-1}])$. From the fact that each $A[d_S^{-1}]/(R_{k-1} \cap A[d_S^{-1}])$ is semisimple the semisimplicity of R_k/R_{k-1} follows. The rest of the Proposition follows from description of the canonical isomorphism (5) recalled above. $\square$

3.4. No broken circuits. In order to finish our analysis of R as a D_T module we have to understand the dimension of V_W for each component W or better yet, to give a basis for these spaces. From now on we shall choose a component W .

Fix a total order on the set of characters $\overline{\Delta}$. Set

$$\Delta_W = \{(a, \chi) \in \Delta \mid \chi \in \Sigma \text{ and } 1 - a\chi = 0, \text{ on } W\}.$$

The projection π (cf. 3.1) of Δ to $\overline{\Delta}$ restricted to Δ_W is injective thus the total ordering on $\overline{\Delta}$ induces a total ordering on Δ_W .

Definition 2. A subset $S = \{(a_1, \chi_1) < \dots < (a_k, \chi_k)\}$ of Δ_W is called a non broken circuit on W (relative to our chosen ordering) if

- (1) S is associated to W

- (2) There is no $(a, \chi) \in \Delta_W$ and no $e = 1, \dots, k$ with the property that $(a, \chi) < (a_e, \chi_e)$ and $\chi, \chi_e, \dots, \chi_k$ are linearly dependent characters.

Theorem 3.5. *The classes $v_{S,W}$, as S runs among the non broken circuits in Δ_W form a basis of V_W .*

Proof. We first show that the classes of $v_{S,W}$ with S a non broken circuit, span V_W . For this let us order the S 's associated to W lexicographically. It is clear that the minimum S in this ordering is non broken.

Pick a general $S = \{(a_1, \chi_1) < \dots < (a_k, \chi_k)\}$ associated to W and assume there is $(a, \chi) \in \Delta_W$ with $\chi, \chi_e, \dots, \chi_k$ linearly dependent characters, and $(a, \chi) < (a_e, \chi_e)$. We can also assume, by induction, that if we remove one of the $\chi_i, e \leq i \leq k$ the remaining elements $\chi, \chi_e, \dots, \check{\chi}_i, \dots, \chi_k$ are linearly independent.

Clearly for each $j = 0, \dots, k-e$, the set $S_j = (S - \{(a_{e+j}, \chi_{e+j})\}) \cup \{(a, \chi)\}$ is associated to W .

Let $\Gamma \supset \Gamma'$ be the sublattices generated by $\chi, \chi_e, \dots, \chi_k$ and $\chi_e, \dots, \chi_k$ respectively. Let U be the torus of coordinate ring $\mathbb{C}[\Gamma]$. We have an associated map $\rho : T \rightarrow U$ under which W maps to the point P_0 of equations $1 - a\chi = 1 - a_e\chi_e = \dots = 1 - a_k\chi_k = 0$.

If we consider only the equations $1 - a_e\chi_e = \dots = 1 - a_k\chi_k = 0$, they define finitely many points $P_0, P_1, \dots, P_t$ in U , we can thus choose an element $\psi \in \mathbb{C}[\Gamma]$ which takes value 1 on P_0 and 0 in the remaining P_i . In particular $\psi = 1$ on W and

$$\phi_{S,W} = \psi \phi_{S,W}, \quad v_{S,W} = \psi v_{S,W}.$$

We have $\psi(1 - a\chi)$ vanishes on all the points $P_0, P_1, \dots, P_t$ hence $\psi(1 - a\chi) = c_e(1 - a_e\chi_e) + \dots + c_k(1 - a_k\chi_k)$, with $c_i \in \mathbb{C}[\Gamma]$, hence:

$$\frac{\psi}{\prod_{i=e}^k (1 - a_i\chi_i)} = \frac{\psi(1 - a\chi)}{(1 - a\chi) \prod_{i=e}^k (1 - a_i\chi_i)} = \sum_{j=e}^k \frac{c_j}{(1 - a\chi) \prod_{i=1, i \neq j}^k (1 - a_i\chi_i)}.$$

Choose a representative $f_{S,W}$ of $\phi_{S,W}$, we have from the previous identity:

$$(6) \quad v_{S,W} = \sum_{j=e}^k \left[\frac{f_{S,W} c_j}{\prod_{i=1}^{e-1} (1 - a_i\chi_i) (1 - a\chi) \prod_{i=e, i \neq j}^k (1 - a_i\chi_i)} \right].$$

We have to understand the elements

$$\left[\frac{f_{S,W} c_j}{\prod_{i=1}^{e-1} (1 - a_i\chi_i) (1 - a\chi) \prod_{i=e, i \neq j}^k (1 - a_i\chi_i)} \right] \in R_k / R_{k-1}.$$

In U , for each j between e and k the equations $1 - a\chi = 0, 1 - a_i\chi_i = 0, \forall e \leq i \leq k, i \neq j$ define a finite set of points U_j . Let $Q := \cup U_j = \{P_0, A_1, \dots, A_f\}$ again a finite set of points including P_0 . We can assume that we have chosen $f_{S,W}$ with the further property of vanishing on all the subvarieties $\rho^{-1}(A_j)$. Consider next the subvariety Z_j of T of equations $1 - a\chi = 0, 1 - a_i\chi_i = 0, \forall 1 \leq i \leq k, i \neq j$, its connected components W_h are elements of the arrangement. Z_j is the preimage under the map $\rho : T \rightarrow U$ of the finite

number of points given by the same equations in U . We claim that on each such W_h the function $f_{S,W}c_j$ is constant. This implies that, modulo the ideal generated by the elements $1 - a\chi = 0, 1 - a_i\chi_i = 0, \forall 1 \leq i \leq k, i \neq j$ the element $f_{S,W}c_j$ is a linear combination of primitive idempotents, hence by (6) $v_{S,W}$ is a linear combination of elements v_{W_h, S_i} with S_i lower than S in the lexicographic order.

In fact this is clear for c_j since this comes from a function on U , as for $f_{S,W}$ we know, by hypothesis that it vanishes on all the components W_h which map under ρ to a point $A_p, p = 1, \dots, f$ thus it only remains to analyze the components lying over P_0 . On these components also $1 - a_j\chi_j$ vanishes and so they are part of the components of the subvariety of equations $1 - a_i\chi_i = 0, 1 \leq i \leq k$. By construction, on this subvariety, $f_{S,W}$ is the characteristic function of W , hence the claim. From this, everything follows by induction.

We now need to show that the vectors $v_{S,W}$ with S a non broken circuit are linearly independent. Take a basis $\xi_1, \dots, \xi_k$ of Σ and complete it to a basis $\xi_1, \dots, \xi_r$ of Λ . Then $I_n = (1 - b_1\xi_1, \dots, 1 - \xi_k) \subset A$ for suitable $b_j \in \mathbb{C}^*$. Consider the completion B of the local ring A_{I_n} at its maximal ideal. We can identify B with the power series ring $K[[x_1, \dots, x_k]]$ with $x_j = \log(b_j\xi_j)$ and $K = \mathbb{C}(\xi_{k+1}, \dots, \xi_r)$, the quotient field of A/I_n . From now on we shall identify A with its image under its canonical inclusion in B . Thus $b_j\xi_j$ equals $\exp(x_j) = \sum_s x_j^s/s!$. Similarly for $(a, \chi) \in \Delta_W$, $a\chi$ equals $\exp(z_\chi)$ where if $\chi = \xi_i^{m_1} \dots \xi_k^{m_k}$, $z_\chi = m_1x_1 + \dots + m_kx_k$, in particular this implies that, in B , we have $1 - a\chi = z_\chi f_\chi$ with f_χ invertible in B and congruent to 1 modulo the maximal ideal. On the other hand, if $(a, \chi) \notin \Delta_W$ then $1 - a\chi$ is invertible in B . It follows that

$$\hat{R} = B\left[\prod_{(a,\chi) \in \Delta} \frac{1}{(1 - a\chi)}\right] = B\left[\prod_{(a,\chi) \in \Delta_W} \frac{1}{z_\chi}\right]$$

Let us define a filtration of $\hat{R}$ in a completely analogous way to the filtration of R . Notice that $R_s \subset \hat{R}_s$ for each s , so that we get a map $j : R_k/R_{k-1} \rightarrow \hat{R}_k/\hat{R}_{k-1}$. The above considerations clearly imply that for any $S = \{(a_1, \chi_1) < \dots < (a_k, \chi_k)\}$ associated to W we have that $j(v_{S,n})$ equals the class w_S of $1/(z_{\chi_1} \dots z_{\chi_k})$. Using this, our claim follows immediately from the linear independence the w_S 's as S runs among the non broken circuits which is well known (see [23]). $\square$

3.5. Partial fractions. Keeping the notations of the previous section consider a component W of the arrangement of codimension k . Its associated character group $\Sigma(W) := \Sigma$ is a split direct summand of Λ of rank k and W is defined by the ideal generated by a maximal ideal $\mathfrak{n}$ in $\mathbb{C}[\Sigma]$. By assumption there exists at least a subset $S \subset \Delta$ associated to W . This means that the irreducible subvariety W is a connected component of the intersection of the $H_{(a,\chi)}$ with $(a, \chi) \in S$.

Take a basis $\xi_1, \dots, \xi_k$ of $\Sigma(W)$ and complete it to a basis $\xi_1, \dots, \xi_r$ of Λ . For each $i = 1, \dots, r$, set $\partial_i = \partial/\partial\xi_i$. This allows us to introduce a commutative subring in D_T namely $D_W := \mathbb{C}[\xi_{k+1}^{\pm 1}, \dots, \xi_r^{\pm 1}][\partial_1, \dots, \partial_k]$.

In R_k/R_{k-1} consider the component $N(W) \otimes V_W$. Let us remark that Lemma 2.1 implies that as a D_W -module $N(W)$ is free of rank 1, generated by the class of 1.

Also remark that by Theorem 3.5, we have a basis of V_W given by the vectors $v_{S,W}$ as S runs through the non broken circuits in Δ_W . We choose a representative $f_{S,W}/\prod_{(a,\chi) \in S}(1-a\chi)$ of $v_{S,W}$ where $f_{S,W}$ is a representative of the primitive idempotent $\phi_{S,W}$. In this way we lift V_W to a subspace $\tilde{V}_W$ lying in the ring obtained from $\mathbb{C}[\Sigma]$ inverting the product of the elements $1-a\chi$ with $(a,\chi) \in \Delta_W$. Let $M(W)$ be the D_W -module generated by $\tilde{V}_W$, $M(W)$ is a free D_W -module with basis any linear basis of $\tilde{V}_W$, in particular the elements $f_{S,W}/\prod_{(a,\chi) \in S}(1-a\chi)$. Under the quotient map $R_k \rightarrow R_k/R_{k-1}$, $M(W)$ maps isomorphically to $N(W) \otimes V_W$. In particular we get,

Corollary 3.6. $R = \oplus_W M(W)$.

If the codimension of W is k , $M(W) \subset R_k$ and π_k maps $M(W)$ isomorphically onto $N(W) \otimes V_W$.

We can make explicit the previous decomposition in terms of an explicit linear basis and a suitable choice of representatives of the primitive idempotents.

For every non broken linearly independent subset $\Gamma = \{\psi_1, \dots, \psi_k\} \subset \Delta$ let us denote by $\Lambda(\Gamma), \overline{\Lambda}(\Gamma)$ the lattice that Γ generates and the set of elements of Λ which are torsion modulo $\Lambda(\Gamma)$. The idempotents $\phi_{\Gamma,W}$ of the ring $\mathbb{C}[\Lambda]/\langle 1-a_1\psi_1, \dots, 1-a_k\psi_k \rangle$ can be also described in terms of characters of the finite group $\overline{\Lambda}(\Gamma)/\Lambda(\Gamma)$. First if all the $a_i = 1$, if $\Phi(\Gamma)$ denotes the character group, given $\lambda \in \Phi(\Gamma)$ a representative ϕ_λ of the corresponding idempotent is constructed as follows, let $h_\Gamma := |\overline{\Lambda}(\Gamma)/\Lambda(\Gamma)|$, take representatives $\xi_1, \dots, \xi_h \in \overline{\Lambda}(\Gamma)$ for the cosets of $\overline{\Lambda}(\Gamma)/\Lambda(\Gamma)$ then:

$$(7) \quad \phi_\lambda = \frac{1}{h_\Gamma} \sum_{i=1}^{h_\Gamma} \lambda(\xi_i^{-1}) \xi_i.$$

We can further normalize the x_i as follows. We think of Λ as being contained in the rational vector space with basis the elements ψ_i then we can uniquely choose the x_i of the form $\prod_j \psi_j^{a_{ij}}$ with $0 \leq a_{ij} < 1$ rational numbers. In this way ϕ_λ is a well defined element of R .

When the $a_i \neq 1$ we must make a change of variables in the ring $\mathbb{C}[\Lambda]$ so that $a_i\psi_i$ becomes one of the group elements (apply an automorphism of translation). We then have

Theorem 3.7. *Every element $f \in R$ can be uniquely expressed in a decomposition in partial fractions:*

$$(8) \quad f = \sum_{\Gamma} \sum_{\lambda \in \Phi(\Gamma)} \sum_{n_1, \dots, n_k} \frac{a_{n_1, \dots, n_k}^{x_i, \Gamma}}{(1 - a_1 \psi_1)^{n_1} \dots (1 - a_k \psi_k)^{n_k}}$$

As x_i runs over a set of representatives of the cosets of $\bar{\Lambda}(\Gamma)/\Lambda(\Gamma)$ or as sum:

$$(9) \quad f = \sum_{\Gamma} \sum_{\lambda \in \Phi(\Gamma)} \sum_{n_1, \dots, n_k} \frac{a_{n_1, \dots, n_k}^{\lambda, \Gamma} \phi_{\lambda}}{(1 - a_1 \psi_1)^{n_1} \dots (1 - a_k \psi_k)^{n_k}}$$

as Γ runs over all the no broken circuit sets and $a_{n_1, \dots, n_k}^{\lambda, \Gamma} \in \mathbb{C}[\xi_{k+1}^{\pm 1}, \dots, \xi_r^{\pm 1}]$ in the corresponding ring.

Proof. The pairs $\Gamma, \lambda \in \Phi(\Gamma)$ are in 1 – 1 correspondence with the components W of the arrangement and we have that the elements $\frac{\xi_{k+1}^{h_1} \dots \xi_r^{h_r} \phi_{\lambda}}{(1 - a_1 \psi_1)^{n_1} \dots (1 - a_k \psi_k)^{n_k}}$, $n_i \in \mathbb{N}$, $h_i \in \mathbb{Z}$ form a basis of the corresponding vector space $M(W)$. Summing these spaces over the components associated to Γ we have a free module over the ring $D_W := \mathbb{C}[\xi_{k+1}^{\pm 1}, \dots, \xi_r^{\pm 1}][\partial_1, \dots, \partial_k]$ with basis the representatives x_i . Then the formula is clear. $\square$

4. COHOMOLOGY

In this section we shall determine the additive structure of the cohomology of the open set $\mathcal{A}_{\Delta} \subset T$ which is the complement of the union of the hypersurfaces of $H_{(a, \chi)}$ of equation $1 - a\chi = 0$ for $(a, \chi) \in \Delta$.

Let $\Xi = \bigwedge(\omega_1, \dots, \omega_r)$ denote the exterior algebra of invariant differential forms, $\omega_i := d \log \xi_i$ for our basis of characters.

Due to the Theorem of Grothendieck [16], the De Rham cohomology of $\mathcal{A}_{\Delta}$ can be computed via its algebraic De Rham complex $\Omega := R \otimes \Xi$.

It is convenient to write the differential in an invariant way.

$$df = \sum_i \frac{\partial f}{\partial \xi_i} d\xi_i = \sum_i \partial_i(f) \omega_i.$$

Using the decomposition of Corollary 3.6, we also have:

$$\Omega = \oplus_W M(W) \otimes \Xi.$$

Next we claim that $M(W) \otimes \Xi$ is a subcomplex.

For this it is enough to see that each $D_W f_{S, W} / \prod_{(a, \chi) \in S} (1 - a\chi) \Xi$ is a complex. Remark that, if $a \in D_W$ we have $\partial_i(a f_{S, W} / \prod_{(a, \chi) \in S} (1 - a\chi)) = a \partial_i f_{S, W} / \prod_{(a, \chi) \in S} (1 - a\chi)$ if $i \leq k$ thus $a \partial_i \in D_W$. Instead, if $i > k$ we have $\partial_i(a) = [\partial_i, a] \in D_W$ and $\partial_i f_{S, W} / \prod_{(a, \chi) \in S} (1 - a\chi) = 0$ and still the claim follows.

Lemma 4.1. *For each component W , the complex $M(W) \otimes \Xi$ is isomorphic to the tensor product of two complexes M_1 and M_2 .*

M_1 is the De Rham complex for the variety W which is isomorphic as an algebraic variety, to a $r - k$ dimensional torus.

$M_2 = \mathbb{C}[\partial_1, \dots, \partial_k] \otimes \bigwedge(\omega_1, \dots, \omega_k) \otimes V_W$, with differential

$$\delta(p \otimes \omega \otimes v) = \sum_{i=1}^k \partial_i p \otimes \omega_i \wedge \omega \otimes v$$

for $p \in \mathbb{C}[\partial_1, \dots, \partial_k]$, $\omega \in \bigwedge(\omega_1, \dots, \omega_k)$, $v \in \tilde{V}_W$.

$$H^s(M_2) = \begin{cases} 0 & \text{if } s < k \\ V_W \otimes \omega_1 \wedge \dots \wedge \omega_k & \text{if } s = k \end{cases}$$

Proof. Using the choices made above, we can identify the coordinate ring of W with $\mathbb{C}[\xi_{k+1}^{\pm 1}, \dots, \xi_r^{\pm 1}]$. Thus $M_1 := \mathbb{C}[\xi_{k+1}^{\pm 1}, \dots, \xi_r^{\pm 1}] \otimes \bigwedge(\omega_{k+1}, \dots, \omega_r)$, with the usual De Rham differential, is the De Rham complex of W .

On the other hand, take $f \otimes \phi_1 \in M_1$, $p \otimes \phi_2 \otimes v \in M_2$ (with the obvious notations). It is immediate to verify that the product

$$f \otimes \phi_1 \otimes p \otimes \phi_2 \otimes v \mapsto fp \otimes \phi_1 \wedge \phi_2 \otimes v$$

induces an isomorphism of complexes $M(W) \otimes \Xi \simeq M_1 \otimes M_2$.

The complex M_2 is the tensor product of the vector space V_W (thought of as trivial complex in degree 0) and the complexes concentrated in degrees 0,1, $\mathbb{C}[\partial_i] \rightarrow \mathbb{C}[\partial_i]\omega_i$ with $d(p) = \delta_i p \omega_i$, whose cohomology is generated by the class of ω_i in degree 1. The statement about the cohomology of M_2 follows immediately from this. $\square$

As an immediate consequence we obtain in a more explicit form the result of Looijenga, [19], §2.4.3.

Theorem 4.2. *For each integer $i \geq 0$, we have a (non canonical) decomposition, as W runs over the components of the arrangement:*

$$H^i(\mathcal{A}_\Delta) = \oplus_W H^{i-\text{codim}W}(W) \otimes V_W.$$

More precisely a basis for the cohomology can be given by the classes of the forms

$$(10) \quad \frac{f_{S,W}}{(1 - a_1 \chi_1) \dots (1 - a_k \chi_k)} d \log \xi_1 \wedge \dots \wedge d \log \xi_k \wedge d \log \xi_{j_1} \wedge \dots \wedge d \log \xi_{j_s}.$$

Here W runs over the set of components, $S = \{(a_1, \chi_1) < \dots < (a_k, \chi_k)\}$ runs over the non broken circuits in Δ_W , and for given (S, W) , $f_{S,W} \in \mathbb{C}[\Sigma]$ is a chosen representative of the primitive idempotent $\phi_{S,W}$. Finally $\xi_1, \dots, \xi_r$ is a basis of the character group such that the χ_i are in the subgroup spanned by the first k elements.

Remark 4.3. 1) *The previous forms do not depend on the choice of a complement of the lattice spanned by $\xi_1, \dots, \xi_k$ in Λ but only on the choice of a basis of $\Lambda / \langle \xi_1, \dots, \xi_k \rangle$.*

2) The filtration of R by the subspaces R_k , induces for each i , a natural filtration $H^i(\mathcal{A}_\Delta)$ on cohomology such the k^{th} graded component is.

$$H^i(\mathcal{A}_\Delta)_k / H^i(\mathcal{A}_\Delta)_{k-1} = \bigoplus_{W | \text{codim } W = k} H^{i-k}(W) \otimes V_W.$$

3) Using Theorem 4.2 one can write down an explicit formula for the Poincaré polynomial of $\mathcal{A}$ in terms of the poset of components. It is easily seen that this formula coincides with the formula given by Looijenga in [19], 2.4.3.

With the previous notations we can also introduce a different basis which will be useful for the future computations:

Proposition 4.4. *The forms*

$$f_{S,W} d\log(1 - a_1\chi_1) \wedge \cdots \wedge d\log(1 - a_k\chi_k) \wedge d\log \xi_{j_1} \wedge \cdots \wedge d\log \xi_{j_s}$$

(where $k < j_1 < \cdots < j_s \leq r$), are closed and their classes give a basis of the cohomology of $\mathcal{A}_\Delta$.

Proof. The fact that these forms are closed follows as in the previous arguments. These forms are obtained by an invertible triangular change of basis with respect to the forms (10). In fact

$$f_{S,W} d\log(1 - a_1\chi_1) \wedge \cdots \wedge d\log(1 - a_k\chi_k) = f_{S,W} \prod_i \frac{-a_i\chi_i}{(1 - a_i\chi_i)} d\log \chi_i.$$

Substitute for $-a_i\chi_i$ the sum $(1 - a_i\chi_i) - 1$ and develop all the terms getting the required triangular expression. $\square$

For our work the top dimension $i = r$ is particularly important. In this case the forms

$$(11) \quad f_{S,W} d\log(1 - a_1\chi_1) \wedge \cdots \wedge d\log(1 - a_k\chi_k) \wedge d\log \xi_{j_1} \wedge \cdots \wedge d\log \xi_{j_{r-k}}$$

are independent of the way in which we complete the basis $\xi_1, \dots, \xi_k$ except for the orientation which we fix in an arbitrary way. Their classes will be denoted by $\omega_{S,W}$. Such a class is said to be *associated* to W .

Given a top cohomology class ψ when we expand ψ in the basis $\omega_{S,W}$ we call the coefficients the *local residues* of ψ at S, W and write:

$$\psi = \sum_{S,W} \text{res}_{S,W}(\psi) \omega_{S,W}.$$

In particular, the form

$$(12) \quad \omega_{\emptyset,T} = d\log \xi_1 \wedge \cdots \wedge d\log \xi_r$$

depends only on the orientation of the basis $\xi_1, \dots, \xi_r$ of Λ and will be denoted by ω_T (or just ω).

4.1. The local residue. Let P be one of the points of the arrangement, thus we have as usual a list Δ_P of the divisors $1 - a_i\psi_i$ passing through P .

Consider $\mathfrak{p}$ the tangent space of T in 1 and let $e_P : t \mapsto P \exp(t)$ which gives local coordinates around P .

The preimage of one of the divisors $1 - a_i\psi_i = 0$ is a hyperplane whose equation h_i is defined by $\psi_i(e^t) = e^{h_i(t)}$. We can restrict to a small enough ball of center 0, so that the preimage of $\mathcal{A}_\Delta$ is exactly the complement of the intersection of these hyperplanes with this ball.

The total ordering on Δ_P transfers to a total ordering of the vectors h_i which preserves the notion of no broken circuits. From a topological point of view this is diffeomorphic to the complement of this hyperplane arrangement. Let us now compute now the pull back under e_P of the top forms and top cohomology classes $\omega_{S,W}$ defined in (11).

Theorem 4.5. 1) Under e_P^* all cohomology classes which are not associated to P go to 0.

2) The cohomology class of the form

$$e_P^*(f_{S,P} d \log(1 - a_1\psi_1) \wedge \dots \wedge (1 - a_r\psi_r))$$

equals that of the form $d \log h_1 \wedge \dots \wedge d \log h_r$.

Proof. Consider one of the forms $d \log(1 - a\psi)$ if $(1 - a\psi)(P) \neq 0$ we can choose (locally) a determination of the logarithm so if the ball is small enough the pull back of this form is exact. The same applies to the forms $d \log \chi$. If instead $(1 - a\psi)(P) = 0$, with $\psi = e^h$ we have that:

$$e_P^* d \log(1 - a\psi) = d \log h + \gamma,$$

with γ holomorphic on the ball.

One concludes with two remarks.

A product of $< r$ factors of type $d \log h_i$ times a holomorphic $r - k$ form is exact.

The form $f(t) d \log h_1 \wedge \dots \wedge d \log h_r$ is cohomologous to $f(0) d \log h_1 \wedge \dots \wedge d \log h_r$. $\square$

It is well known [23], that the forms $w_S := d \log h_1 \wedge \dots \wedge d \log h_r$ as $S := \{h_1, \dots, h_r\}$ runs over the no broken circuits of the hyperplane arrangement, form a basis of the cohomology of the complement of the hyperplanes. We get

Theorem 4.6. Given a point P of the arrangement and a differential form of top degree Ψ we have

$$(13) \quad e_P^*([\Psi]) = \sum_{S \in \Delta_P} \text{res}_{S,P}([\Psi]) w_S.$$

where $[\Psi]$ denotes the cohomology class of Ψ .

In [9] we have constructed geometrically a dual homology basis, and shown how one can explicitly compute the integral of the form on each element of

the basis as a local residue around a new point at *infinity* in a suitable blow up which transform the divisor of hyperplanes into one with normal crossing. This then becomes a method to compute the local residues $res_{S,P}([\Psi])$.

In fact we will need to analyze, for suitable Ψ , the function

$$c(\beta) := res_{S,P}([\beta^{-1}\Psi])$$

as $\beta \in \Lambda$. Let us briefly discuss an important qualitative feature of this number as a function of β .

Definition 3. *A function g , on Λ will be called a **periodic polynomial** if it is a polynomial on each coset of some sublattice Λ' of Λ .*

Remark that as sublattice we can take $n\Lambda$ for some positive integer n . If f is a periodic polynomial on Λ , it is also a periodic polynomial on $1/n\Lambda$ for every positive integer n .

Proposition 4.7. *$c(\beta) = res_{S,P}([\beta^{-1}\Psi])$ is a periodic polynomial on Λ .*

Proof. In the coordinates $y_1, \dots, y_r$ of the blow up (see [9] for the explicit formulas), the form Ψ develops as:

$$\Psi = \frac{1}{y_1^{h_1} \dots y_r^{h_r}} G(y_1, \dots, y_r) dy_1 \wedge \dots \wedge dy_r$$

where G is a holomorphic function near zero. As for β^{-1} it can be written as $\beta^{-1}(p)$ times a convergent power series in the y 's with coefficient which depend polynomially on β . From this the claim follows since $\beta^{-1}(p)$ is constant on the sublattice Λ' spanned by $\psi_1, \dots, \psi_r$. $\square$

In the way we have organized our discussion, the computation of the local residues is performed by viewing the toric arrangement locally, around each of the points P of the arrangement. We have used the fact that locally the arrangement is isomorphic to a hyperplane arrangement, to which we then apply our geometric theory of blow ups. In fact, although we do not use this explicitly, the theory is even global. It is clear that the stratification of a torus induced by the toric arrangement is *conical* as defined in [22]. The theory of irreducible strata and nested sets can be computed locally and thus reduced to the hyperplane case where it is well understood. Then, by applying the theory developed in [22], we can construct a smooth model $\tilde{T}$ with a proper map $\pi : \tilde{T} \rightarrow T$. π is isomorphic on the complement $\mathcal{A}_\Delta$ of the arrangement. The divisor in $\tilde{T}$, complement of $\mathcal{A}_\Delta$, has normal crossings and it has globally the combinatorics described locally, for all the points P of the arrangement, by the corresponding hyperplane arrangements.

4.2. Coverings. Let us now consider a finite n -sheeted covering $\pi : U \rightarrow T$ of tori. It corresponds to an embedding $\Lambda \subset M$ of character groups, with Λ of index n in M . It is a Galois covering with Galois group the dual $(M/\Lambda)^*$ of M/Λ .

The set Δ clearly defines also an arrangement in U , to every component W of the arrangement in T correspond the connected components of $\pi^{-1}(W)$.

Let us denote by $\mathcal{A}_\Delta^T, \mathcal{A}_\Delta^U$ the complement for the two arrangements, we have thus a map $\pi : \mathcal{A}_\Delta^U \rightarrow \mathcal{A}_\Delta^T$ and in cohomology $\pi^* : H^*(\mathcal{A}_\Delta^T) \rightarrow H^*(\mathcal{A}_\Delta^U)$.

The vector spaces of invariant derivations for T and U coincide. Thus we can take a common basis $\partial_1, \dots, \partial_r$ and we have

$$(14) \quad D_U = \mathbb{C}[U][\partial_1, \dots, \partial_r] = \mathbb{C}[U] \otimes_{\mathbb{C}[T]} D_T.$$

For a D_T module M we can construct the *induced module* on U as:

$$D_U \otimes_{D_T} M = \mathbb{C}[U] \otimes_{\mathbb{C}[T]} M.$$

The action of a derivation ∂_i on $a \otimes m$, $a \in \mathbb{C}[U]$ is clearly:

$$\partial_i(a \otimes m) = \partial_i(a) \otimes m + a \otimes \partial_i(m).$$

The following Proposition, describing the induction of the modules $N(W)$ of §2.2, is only a reformulation of Lemma 2.2:

Proposition 4.8. *Given a covering $\pi : U \rightarrow T$ and an irreducible component $W \subset T$ of an arrangement, there is a canonical isomorphism of D_U modules*

$$\mathbb{C}[U] \otimes_{\mathbb{C}[T]} N(W) \simeq \bigoplus_{i=1}^h N(W_i).$$

Where the W_i are the irreducible components of $\pi^{-1}(W)$.

This has an important consequence for the canonical filtration of $\mathbb{C}[U]_\Delta := \mathbb{C}[U][d^{-1}]$. Start from the fact that $\mathbb{C}[U]_\Delta = \mathbb{C}[U] \otimes_{\mathbb{C}[T]} \mathbb{C}[T]_\Delta$.

Theorem 4.9. *For each $k = 0, \dots, r$:*

$$(\mathbb{C}[U]_\Delta)_k = \mathbb{C}[U] \otimes_{\mathbb{C}[T]} (\mathbb{C}[T]_\Delta)_k.$$

$$(\mathbb{C}[U]_\Delta)_k / (\mathbb{C}[U]_\Delta)_{k-1} = \mathbb{C}[U] \otimes_{\mathbb{C}[T]} [(\mathbb{C}[T]_\Delta)_k / (\mathbb{C}[T]_\Delta)_{k-1}]$$

hence we have canonical isomorphisms:

$$(\mathbb{C}[U]_\Delta)_k / (\mathbb{C}[U]_\Delta)_{k-1} \simeq \bigoplus_W \mathbb{C}[U] \otimes_{\mathbb{C}[T]} N_T(W) \otimes V_W \simeq \bigoplus_W \bigoplus_Z N_U(Z) \otimes V_W$$

W runs over the components of the arrangement in T of codimension k and, for given W , Z runs over the components of the arrangement in U which lie over W .

Proof. Since $\mathbb{C}[U]$ is a free module over $\mathbb{C}[T]$ the functor $\mathbb{C}[U] \otimes_{\mathbb{C}[T]} -$ is exact hence the claims follow since the k -th level of the filtration of $\mathbb{C}[U]_\Delta$ can be characterized as the maximal D_U submodule whose characteristic variety is the union of conormal bundles of subvarieties of codimension $\leq k$. $\square$

Since $\Lambda_{\mathbb{R}} = M_{\mathbb{R}}$ we fix once and for all an orientation of this vector space which induces compatible orientations for all coverings. Then we have the canonical classes ω_T, ω_U as in formula 12 and:

Lemma 4.10. *If π is of degree n :*

$$\pi^*(\omega_T) = n\omega_U$$

Proof. By the theory of elementary divisors, there is an oriented basis $\mu_1, \dots, \mu_r$ of M and positive integers $n_1, \dots, n_r$ such that $n = \prod_i n_i$ and $\mu_1^{n_1}, \dots, \mu_r^{n_r}$ is a basis of Λ . Using these two bases the claim is clear. $\square$

Let us compute in general $\pi^*(\omega_{S,W})$ where $\omega_{S,W}$ is defined in formula (11).

Given the lattice $\langle S \rangle \subset \Lambda$ its associated $\overline{\langle S \rangle}$ of elements of Λ which are torsion modulo $\langle S \rangle$ and finally $\langle \tilde{S} \rangle$ of elements of M which are torsion modulo $\langle S \rangle$ we can construct summands:

$$\Lambda = \overline{\langle S \rangle} \oplus \Pi, \quad M = \langle \tilde{S} \rangle \oplus \tilde{\Pi}, \quad \Pi \subset \tilde{\Pi}$$

We have that $\pi^{-1}(W)$ has $[\langle \tilde{S} \rangle : \overline{\langle S \rangle}]$ components while, if $\xi_1, \dots, \xi_{r-k}$ resp. $\eta_1, \dots, \eta_{r-k}$ are bases of Π resp. of $\tilde{\Pi}$ with the same orientation, we have

$$(15) \quad d \log \eta_1 \wedge \dots \wedge d \log \eta_{r-k} = p^{-1} d \log \xi_1 \wedge \dots \wedge d \log \xi_{r-k}, \quad p = [\tilde{\Pi} : \Pi].$$

For every component W' of $\pi^{-1}(W)$ we can clearly choose a representative, $f_{S,W'}$ of the corresponding primitive idempotent in such a way that

$$f_{S,W} = \sum_{W' \text{ component of } \pi^{-1}(W)} f_{S,W'}.$$

We then set

$$\omega_{S,W'} := f_{S,W'} d \log(1 - a_1 \chi_1) \wedge \dots \wedge d \log(1 - a_k \chi_k) \wedge d \log \eta_1 \wedge \dots \wedge d \log \eta_{r-k}$$

Proposition 4.11. *We have that:*

$$\pi^*(\omega_{S,W}) = p^{-1} \sum_{W'} \omega_{S,W'}$$

as W' runs over the components of $\pi^{-1}(W)$.

Proof. Everything follows from identity (15). $\square$

Notice that in the case in which W is a point we have that $p = 1$.

Let us specialize to one important case. Let U be the torus with character group $M = \frac{1}{m}\Lambda$ for some integer m so that the degree of the map $\pi : U \rightarrow T$ is m^r .

Inverting the elements of Δ is equivalent to inverting the elements of:

$$\Delta^{1/m} := \{(b, \sigma), \quad b \in \mathbb{C}^*, \quad \sigma \in \frac{1}{m}\Lambda, \quad | (b^m, \sigma^m) \in \Delta \}.$$

In fact let $(a, \chi) \in \Delta$, choose b with $b^m = a$. We have m pairs $(\zeta^i b, \sigma)$, $i = 0, \dots, m-1$, $\zeta = e^{2\pi i/m}$ with $((\zeta^i b)^m, \sigma^m) = (a, \chi)$ where σ is uniquely determined. Moreover

$$1 - a\chi = \prod_{i=0, \dots, m-1} (1 - b\zeta^i \sigma).$$

Take a component $W \subset T$ of the arrangement $\mathcal{R}_\Delta$. This is a connected component of the variety of equations $a_i \chi_i = 1$ $i = 1, \dots, k$, where $(a_i, \chi_i) \in \Delta$ and the χ_i are linearly independent.

$\pi^{-1}(W)$ consists of m^k connected components, one component for each of the varieties $\zeta^j b_i \sigma_i = 1$, $i = 1, \dots, k$ as $b_i^m = a_i$, $\sigma^m = \chi_i$ and the exponents j vary in the m^k possible different ways.

Recall that $\overline{\Delta}$ has been defined as the set of characters χ such that there is $a \in \mathbb{C}^*$ with $(a, \chi) \in \Delta$ and define $\overline{\Delta}^{1/m}$ in the same way. Notice that $\overline{\Delta}^{1/m}$ is in canonical bijection with $\overline{\Delta}$. Thus the chosen ordering on $\overline{\Delta}$ gives an ordering on $\overline{\Delta}^{1/m}$.

For the given component W , select a component W' of $\pi^{-1}(W)$.

It is then immediate to see that the map $(b, \sigma) \mapsto (b^m, \sigma^m)$ defines a bijection between $\Delta_{W'}^{1/m}$ and Δ_W . This establishes also a bijection between the no broken circuits on W' and the ones on W .

As for cohomology, take a basic form

$$f_{S,W} d \log(1 - a_1 \chi_1) \wedge \dots \wedge d \log(1 - a_k \chi_k) \wedge d \log \xi_{j_1} \wedge \dots \wedge d \log \xi_{j_s}.$$

Choose as complement of the lattice associated the elements χ_i or their m -th roots, in the character lattice of the covering, the one generated by the m - roots η_i of the ξ_i .

Since

$$d \log(1 - a_i \chi_i) = d \log\left(\prod_{j=0}^{m-1} (1 - \zeta^j b_i \sigma_i)\right) = \sum_{j=0}^{m-1} d \log(1 - \zeta^j b_i \sigma_i),$$

$$d \log \xi_i = d \log(\eta_i^m) = m d \log \eta_i,$$

we have that the previous form is a sum of terms

(16)

$$m^s f_{S,W} d \log(1 - \zeta^{j_1} b_1 \sigma_1) \wedge \dots \wedge d \log(1 - \zeta^{j_k} b_k \sigma_k) \wedge d \log \eta_{j_1} \wedge \dots \wedge d \log \eta_{j_s}$$

The function $f_{S,W}$ has value 1 on the preimage in U of W and value 0 on the preimage of all the other components of the variety of equations $1 - a_i \chi_i = 0$.

Thus, with the previous notations, for $S_{\underline{j}} := \{1 - \zeta^{j_1} b_1 \sigma_1, \dots, 1 - \zeta^{j_k} b_k \sigma_k\}$ we have that:

Lemma 4.12. *We can choose representatives $f_{S_{\underline{j}}}$ as W_t runs over all the components of the variety of equations $1 - \zeta^{j_1} b_1 \sigma_1 = \dots = 1 - \zeta^{j_k} b_k \sigma_k = 0$ in U , which lie over W so that:*

$$f_{S,W} = \sum_t f_{S_{\underline{j}}, W_t}$$

Proof. We choose all except one which we define from the previous formula and verify that it is indeed a required representative. $\square$

Remark we should compare the two expressions associated to a component in U using either Δ or $\Delta^{1/m}$.

We have

$$d \log(1 - \zeta^{j_1} b_1 \sigma_1) \wedge \dots \wedge d \log(1 - \zeta^{j_k} b_k \sigma_k) =$$

$$\prod_{i=1}^k \prod_{j=0, j \neq j_i}^{m-1} (1 - \zeta^j b_k \sigma_k) d\log(1 - a_1 \psi_1) \wedge \cdots \wedge d\log(1 - a_k \psi_k)$$

A special case will be needed, when $W = P$ reduces to a point. The 0-dimensional components of the arrangement play a special role in the final formulas. For such P in formula (16) we have $s = 0$ and thus:

$$(17) \quad f_{S,P} d\log(1 - a_1 \chi_1) \wedge \cdots \wedge d\log(1 - a_r \chi_r) = \sum_{Q \in \pi^{-1}(P)} f_{S_{\underline{j}}, Q} d\log(1 - \zeta^{j_1} b_1 \sigma_1) \wedge \cdots \wedge d\log(1 - \zeta^{j_r} b_r \sigma_r).$$

In the right hand side of this formula the components lying over $W = P$ are just the m^r points in $Q \in \pi^{-1}(P)$ the fiber. Each such point Q determines a unique no broken circuit $S_{\underline{j}}$ obtained as m -th root of S . The formula induces a corresponding formula in cohomology.

5. THE COHOMOLOGY ALGEBRA

5.1. A basic identity. We start from a basic formal identity:

$$(18) \quad 1 - \prod_{i=1}^n x_i = \sum_{I \subsetneq \{1, 2, \dots, n\}} \prod_{i \in I} x_i \prod_{j \notin I} (1 - x_j).$$

The proof is by induction on n .

We split the sum in 3 terms: $I = \{1, \dots, n-1\}$, $I \subsetneq \{1, \dots, n-1\}$ and finally $n \in I$. We get

$$\prod_{i=1}^{n-1} x_i (1 - x_n) + (1 - \prod_{i=1}^{n-1} x_i) (1 - x_n) + x_n (1 - \prod_{i=1}^{n-1} x_i) = 1 - \prod_{i=1}^n x_i.$$

Formula (18) implies

$$\sum_{I \subsetneq \{1, 2, \dots, n\}} \frac{1}{(1 - \prod_{i=1}^n x_i)} \prod_{i \in I} \frac{x_i}{(1 - x_i)} = \frac{1}{\prod_{i=1}^n (1 - x_i)}.$$

We want to interpret this formula as an identity between certain differential forms.

Set, for $i = 1, \dots, n$, $\omega_i := d\log(1 - x_i)$, $\psi_i := d\log x_i$. Also for $s = 0, \dots, n$, set $\theta^{(s)} = d\log(1 - \prod_{i=1}^s x_i^{-1} \prod_{j=s+1}^n x_j)$. If we take a proper subset $I = \{i_1 < \dots < i_t\}$ in $\{1, \dots, n\}$ and let $J = \{j_1 < \dots < j_{n-t}\}$ be its complement, we can then define the n -forms:

$$\Phi_I^{(s)} = (-1)^{s_I} \omega_{i_1} \wedge \cdots \wedge \omega_{i_t} \wedge \psi_{j_1} \wedge \cdots \wedge \psi_{j_{n-t-1}} \wedge \theta^{(s)}$$

and

$$\Psi_I = (-1)^{s_I} \omega_{i_1} \wedge \cdots \wedge \omega_{i_t} \wedge \psi_{j_1} \wedge \cdots \wedge \psi_{j_{n-t}}$$

with s_I equal to the parity of the permutation $(i_1, \dots, i_t, j_1, \dots, j_{n-t})$.

We have

Proposition 5.1. *For each $0 \leq s \leq n$, the n -form $\omega_1 \wedge \cdots \wedge \omega_n$ can be written as a linear combination with integer coefficients of the n -forms $\Phi_I^{(s)}$ and Ψ_I .*

Proof. We first deal with the case $s = 0$. In this case, computing we get

$$(19) \quad \Phi_I^{(0)} = (-1)^{|I|} \prod_{i \in I} \frac{x_i}{(1-x_i)} \frac{1}{(1 - \prod_{i=1}^n x_i)} dx_1 \wedge dx_2 \wedge \cdots \wedge dx_n.$$

Thus our identity (18) can be translated into:

$$(20) \quad \sum_{I \subsetneq \{1,2,\dots,n\}} (-1)^{|I|+n} \Phi_I^{(0)} = \omega_1 \wedge \cdots \wedge \omega_n.$$

proving our claim.

In the general case let us observe that

$$d \log(1 - x^{-1}) = d \log \frac{x-1}{x} = -d \log(x) - d \log(1-x).$$

Therefore, the substitution of ix_i with x_i^{-1} for $i = 1, \dots, s$, corresponds in the formula (20) to substitute ω_i with $-\omega_i - \psi_i$ getting a new formula which we can denote as (20'). $\square$

5.2. Formality. In this section we shall assume that our set Δ is *unimodular*, with the terminology and notations of §3.1, this means that given a subset $\Psi \subset \Delta$ which is linearly independent, $\pi(\Psi)$ generates a lattice which is a direct factor in Λ . In particular, if ξ_0 is dependent on $\xi_1, \dots, \xi_k$ with all these elements in Δ we must have that ξ_0 is a product of the ξ_i with exponents ± 1 . We call this a *simple dependency relation*. More generally we say that the element (a_0, ξ_0) is dependent on $(a_1, \xi_1), \dots, (a_k, \xi_k)$ if ξ_0 is a monomial in the ξ_i , $i = 1, \dots, k$ and a_0 is the value of the same monomial in the a_i .

Let us start with a formal construction. Define an exterior algebra in the following generators:

A generator $\lambda_{(a,\chi)}$ for every $(a, \chi) \in \Delta$ and a generator μ_χ for every character $\chi \in \Lambda$.

We then impose a set of relations. Of this first we have the essentially trivial relations:

- (1) $\mu_{\chi_1 \chi_2} = \mu_{\chi_1} + \mu_{\chi_2}$, for all $\chi_1, \chi_2 \in \Lambda$.
- (2) If $\chi_1, \dots, \chi_s$ are dependent the product of elements σ_i where $\sigma_i = \mu_{\chi_i}$ or $\sigma_i = \lambda_{(a_i, \chi_i)}$ is 0.

We come now to the main relations, which generalize the ones for hyperplanes.

Given $(a_0, \chi_0), (a_1, \chi_1), \dots, (a_n, \chi_n) \in \Delta$ with

$$a_0 \chi_0 = \prod_{i=1}^s (a_i \chi_i)^{-1} \prod_{j=s+1}^n a_j \chi_j.$$

We take the formula (20') and substitute ψ_i with μ_{χ_i} and ω_i with λ_{a_i, χ_i} . We get then a formal expression which we impose as a new relation.

We do this for all the simple dependency relation and we call $\mathcal{H}$ the resulting quotient algebra.

Consider the subalgebra H in $\Omega = R \otimes \Xi$ generated by the 1-forms $d \log(1 - a\chi), d \log \chi$. H clearly consists of closed forms so that we obtain a algebra homomorphism

$$f : H \rightarrow H^*(\mathcal{A}_\Delta).$$

It is also easy to verify, using Proposition 5.1, that we have a homomorphism $g : \mathcal{H} \rightarrow H$ given by:

$$g(\lambda_{(a, \chi)}) := d \log(1 - a\chi), \quad g(\mu_\chi) := d \log \chi.$$

Theorem 5.2. *The homomorphisms g, f are isomorphisms.*

Proof. The assumption that Δ is unimodular clearly implies that for each component W and for any S associated to W , the defining ideal of W is generated by the elements $1 - a\chi$ with $(a, \chi) \in S$. In particular S is associated only to one component and thus the idempotent $\phi_{S, W} = 1$. From this and Remark 4.3 we deduce that f is surjective.

The fact that g is surjective is clear from its definition. So, in order to prove our claim, it suffices to see that fg is injective.

We fix a total ordering on Δ in order to apply the theory of no broken circuits. The first set of relations implies that the subalgebra generated by the elements μ_χ is a homomorphic image of the exterior algebra Ξ . So we can consider $\mathcal{H}$ as a Ξ -module.

For a fixed component W we define the subspace $\mathcal{V}_W \subset \mathcal{H}$ spanned by the monomials $\lambda_S = \prod_{(a, \chi) \in S} \lambda_{(a, \chi)}$ (here the product is taken according to the fixed ordering of Δ) as S runs among the subsets of Δ associated to W . The first set of relations also implies that $\mathcal{V}_W$ is annihilated by the elements μ_χ if the character χ is constant on W .

Given a monomial m in the generators $\lambda_{(a, \chi)}$ and μ_{x_i} , we define its weight as the number of factors of m of the first type $\lambda_{(a, \chi)}$. Notice that the weight of $\lambda_S \in \mathcal{V}_W$ equals the codimension of W . Then, whenever S is a broken circuit, the relations of the second type allow to replace it by a product of elements with lower weight or lower in the lexicographical order. This implies that modulo elements of smaller weight any element in $\mathcal{V}_W$ can be written as a linear combination of elements λ_S with S a non broken circuit associated to W .

From this our claim follows immediately from Proposition 4.4. $\square$

Remark 5.3. *We wish to point out that our result shows in particular that in $H^*(\mathcal{A}_\Delta)$ the algebraic relations between the generating forms $d \log(1 - a\chi)$ and $d \log \xi$ resemble, but are more complicated, than the relations of Orlik-Solomon in the case of hyperplane arrangements.*

6. RESIDUES

The rest of this paper is devoted to show the relationship between our work and the theory of counting integer points in polytopes developed by Brion, Szenes, Vergne ([4],[26]). In a way, it is mostly a reformulation in our language of their results and does not claim to be particularly original.

Our main point is that we want to give a geometric interpretation of the sum over certain roots of 1 which appears in their formulas. In our approach this has a clear geometric explanation, first as sum over the zero dimensional components of the toric arrangement and finally as a further sum over suitable *points at infinity* in the blown up model where the arrangement is made of divisors with normal crossing. In this sense we recover a *residue formula* as in the 1-dimensional case. The points at infinity which contribute with a non zero residue depend on the character we choose (in the partition function) in a way which is encoded in the notion of Jeffrey–Kirwan residue.

From now on we shall assume that there is a one parameter group $j : G_m \rightarrow T$ such that for each $(a, \gamma) \in \Delta$, $\langle x_i | j(t) \rangle = t^{m_i}$ with $m_i > 0$.

This assumption implies the following geometric considerations.

Let $\Lambda_{\mathbb{R}} := \Lambda \otimes \mathbb{R}$, the real vector space spanned by the characters. In $\Lambda_{\mathbb{R}}$ denote by $C(\Delta)$, the convex cone of non negative linear combinations of the elements γ for all $(a, \gamma) \in \Delta$. The previous hypothesis implies that $C(\Delta)$ is a *pointed cone*, that is it lies entirely on one side of some hyperplane.

We can thus consider the ring P of formal power series in the characters $\chi \in C(\Delta)$ and its localization Q obtained inverting all characters.

Let us denote the De Rham complex for Q by $\Omega^*(Q)$. Using the form $\omega := \omega_T = d \log \xi_1 \wedge \dots \wedge d \log \xi_r$ where ξ_i is an oriented basis of Λ , we have that the top forms can be identified to $Q\omega$.

Lemma 6.1. $H^r(\Omega^*(Q)) = \mathbb{C}$ and it is generated by the class of ω .

Proof. Notice that, setting as before $\partial_i = \xi_i \frac{\partial}{\partial \xi_i}$, we have that, if $\chi = \prod_i \xi_i^{n_i}$, $\partial_i \chi = n_i \chi$ so that the characters are eigenvectors for the ∂_i . The differential has the form $\psi \mapsto \sum_i \partial_i \psi \wedge d \log \xi_i$ we claim that the exact forms of degree r are exactly the ones which can be written as $f\omega$ where f is a series in the characters without constant term. In one way it is clear, a form $\sum_i \partial_i \psi \wedge d \log \xi_i$ has no constant term, conversely let $f \in Q$ write $f = f_1 + g$ where $f_1 = \sum_{h=-N, h \neq 0}^{\infty} \xi_1^h a_h$ is the sum of all terms containing ξ_1 with a non 0 exponent. Then

$$f_1 \omega = d \sum_{h=-N, h \neq 0}^{\infty} h^{-1} \xi_1^h a_h d \log \xi_2 \wedge d \log \xi_3 \dots \wedge d \log \xi_r$$

We then continue in the same way by induction with g . From this everything follows. $\square$

Definition 4. The canonical map $\Omega^r(Q) \mapsto H^r(\Omega^*(Q)) = \mathbb{C}$ will be called the **residue** and denoted by res .

Notice that, given a function $f \in Q$ we have that $\text{res}(f) := \text{res}(f\omega)$ is the constant term of the Laurent series f . We deduce that, given $f = \sum_{\chi \in \Lambda} a_\chi \chi$ in Q :

$$(21) \quad a_\chi = \text{res}(\chi^{-1}f) = \text{res}(\chi^{-1}f\omega).$$

We have also a canonical map $\Omega^r(R) \mapsto H^r(\Omega^*(R)) = H^r(\mathcal{A}_\Delta)$. This will be called the *total residue* and denoted by $Tres$. Given a form $f\omega \in \Omega^r(R)$ sometimes we shall write just $Tres(f) := Tres(f\omega)$.

Setting

$$i\left(\frac{1}{1-a_\chi}\right) = \sum_{n \geq 0} (a_\chi)^n$$

for each $(a, \chi) \in \Delta$, we get an inclusion morphism $i : R \mapsto Q$.

In the next paragraph it will be convenient to write the lattice Λ additively and write the characters as e^α . Then we can interpret the map $i : R \mapsto Q$ as a map from R to a space of functions on Λ and call it the *Discrete Inverse Laplace Transform*, or DILT for short.

The map i extends to a morphism of De Rham complexes $\Omega^*(R) \mapsto \Omega^*(Q)$ and thus of cohomology groups, which by abuse of notation, we shall still denote by i . In top cohomology this gives the basic formula

$$\text{res}(i(f)) = i(Tres(f)).$$

6.1. Coverings. We pass now to analyze this picture for a given covering $\pi : S \rightarrow T$, of degree n , with Galois group $G = \ker(\pi)$. We then define the algebras, R, P, Q for the corresponding tori and denote them by $R_S, R_T, P_S, P_T, Q_S, Q_T$. The covering corresponds to an inclusion $\Lambda \subset M$ of character groups,. Since $\Lambda_{\mathbb{R}} = M_{\mathbb{R}}$ we fix once and for all an orientation of this vector space which induces compatible orientations for all coverings. Recall the basic formula $\pi^*(\omega_T) = n\omega_S$ for the canonical classes ω_T, ω_S .

We then have commutative diagrams:

$$\begin{array}{ccccc} R_T & \xrightarrow{i} & Q_T & & \Omega^r(R_T) \xrightarrow{Tres} H^r(\Omega^*(R_T)) \xrightarrow{i} H^r(\Omega^*(Q_T)) \\ \pi^* \downarrow & & \pi^* \downarrow & , & \pi^* \downarrow \qquad \qquad \qquad \pi^* \downarrow \qquad \qquad \qquad \pi^* \downarrow \\ R_S & \xrightarrow{i} & Q_S & & \Omega^r(R_S) \xrightarrow{Tres} H^r(\Omega^*(R_S)) \xrightarrow{i} H^r(\Omega^*(Q_S)) \end{array}$$

Under identification of $H^r(\Omega^*(Q_T)), H^r(\Omega^*(Q_S))$ with $\mathbb{C}$ we have the commutative diagram:

$$(22) \quad \begin{array}{ccc} H^r(\Omega^*(Q_T)) & \equiv & \mathbb{C} \\ \pi^* \downarrow & & n \downarrow \\ H^r(\Omega^*(Q_S)) & \equiv & \mathbb{C} \end{array}$$

Let us go back to the special important case of S the torus with character group $M = \frac{1}{m}\Lambda$ for some integer m so that the degree of the map $\pi : S \rightarrow T$ is m^r .

Recall formula (17) defining top forms

$$f_{S,P} d\log(1 - a_1\chi_1) \wedge \cdots \wedge d\log(1 - a_r\chi_r) = \sum_{Q \in \pi^{-1}(P)} f_{S_{\underline{j}},Q} d\log(1 - \zeta^{j_1}b_1\sigma_1) \wedge \cdots \wedge d\log(1 - \zeta^{j_r}b_r\sigma_r).$$

If $\pi^*(S)$ denotes the set of no broken circuits $S_{\underline{j}} = (\zeta^{h_1}b_1, \sigma_1), \dots, (\zeta^{h_r}b_r, \sigma_r)$, obtained as m^{th} roots of S , we have seen that they are in 1-1 correspondence with the points of the fiber $\pi^{-1}(P)$. We shall denote by $\omega_{S,P}, \omega_{S_{\underline{j}},Q}$ the cohomology classes of these forms. Thus we have that:

$$(23) \quad \pi^*(\omega_{S,P}) = \sum_{Q \in \pi^{-1}(P)} \omega_{S_{\underline{j}},Q} = \sum_{S_{\underline{j}} \in \pi^*(S)} \omega_{S_{\underline{j}},Q}.$$

7. THE COUNTING FORMULAS

As we discussed in the introduction, one of the aims of this paper is to understand in a more algebraic geometric language, the counting formulas of [5], [4], [26] for integral points in polytopes.

In this setting we are given a family B of vectors in a lattice Λ (which we treat additively) and, for $\beta \in \Lambda$ we want to count the number c_β of solutions of the equation:

$$\sum_{\alpha \in B} n_\alpha \alpha = \beta, \quad \alpha \in \mathbb{N}.$$

This is clearly the value of a *partition function*, or in other words, the number of integral points of the polytope

$$\Pi_\beta := \{(x_\alpha) \mid \sum_{\alpha \in B} x_\alpha \alpha = \beta, \quad 0 \leq x_\alpha \in \mathbb{R}\}.$$

Of course using the multiplicative notation $\chi_\alpha := e^\alpha$ we have that the numbers c_β are the coefficients of the generating series:

$$(24) \quad f_B := \sum_{\beta} c_\beta e^\beta = \prod_{\alpha \in B} \frac{1}{1 - e^\alpha} = \prod_{\alpha \in B} \frac{1}{1 - \chi_\alpha}$$

Thus the problem is to understand an *inversion formula*, the formula which computes these numbers c_β from f_B .

This is a special case, when $\Delta = \{(1, e^\alpha)\}$, $\alpha \in B$, of our theory. We have seen, for a general Δ , that, for every function $f \in Q$ these numbers are computed as residues by formula (21).

Our next goal is to give, for any function

$$f := \frac{1}{\prod_{(a,\chi) \in \Delta} (1 - a\chi)^{h_{a,\chi}}}, \quad h_{a,\chi} \in \mathbb{N}$$

and a character $\beta \in C(\Delta)$ a computable expression of $\text{res}(\beta^{-1}f)$ in terms of a set (depending on β) of local residues at the points of the arrangement. This more general setting gives a weighted version of the partition function which is essentially a variation of the so called Euler–MacLaurin sums of [5],[4].

7.1. General partial fractions. The results we present from now on are just simple reformulations of the results by Szenes–Vergne [26]. We only stress more the role of no–broken circuits which we feel may improve the algorithmic nature of these formulas.

As before, let us take a finite subset $\Delta \subset \mathbb{C}^* \times \Gamma$. We make two assumptions on Δ :

- (1) The projection $\pi : \Delta \rightarrow \Gamma$ gives pairwise non proportional vectors.
- (2) The characters χ with as (a, χ) varies in Δ , span Γ .

Consider now a pair (b, ξ) with $b \in \mathbb{C}^*$ and $\xi \in \Gamma_{\mathbb{Q}} = \Gamma \otimes \mathbb{Q}$.

We say that (b, ξ) is compatible with Δ if there is a positive integer k such that $(b^k, \xi^k) \in \Delta$. Notice that by assumption (1), we have that the pair (b^k, ξ^k) is uniquely determined by the pair (b, ξ) and will be called the element of Δ corresponding to (b, ξ) .

We now fix once and for all a total ordering on Δ . This ordering induces in an obvious way a partial ordering on the set of pairs compatible with Δ . We shall consider sequences, possibly with repetitions, $(b_1, \xi_1), \dots, (b_r, \xi_r)$, of elements compatible with Δ and we shall always assume that for each $i = 1, \dots, r-1$, either $(b_i, \chi_i) \leq (b_{i+1}, \chi_{i+1})$ or (b_i, χ_i) and (b_{i+1}, χ_{i+1}) are not compatible.

Definition 5. *A non broken circuit is:*

a sequence $(b_1, \xi_1), \dots, (b_r, \xi_r)$, of distinct elements compatible with Δ , satisfying the following two conditions:

- (1) *For each $i = 1, \dots, r$ there is not a pair $(a, \chi) \in \Delta$ with $(a, \chi) < (b_i, \chi_i)$ and a sequence of non zero integers $m, n_i, \dots, n_r$ such that:*

$$(a\chi)^m (b_i \xi_i)^{n_i} \dots (b_r \xi_r)^{n_r} = 1.$$

- (2) *$\xi_1, \dots, \xi_r$ are linearly independent.*

We start with another basic identity similar to the one given in 5.1.

Lemma 7.1.

$$(25) \quad 1 - \prod_{i=1}^r z_i = \sum_{\emptyset \subsetneq I \subset \{1, \dots, r\}} \prod_{i \in I} (-1)^{|I|+1} (1 - z_i)$$

Proof. The proof is by induction on r . The case $r = 1$ is clear. In general, using the inductive hypothesis, we have

$$\sum_{\emptyset \subsetneq I \subset \{1, \dots, r\}} \prod_{i \in I} (-1)^{|I|+1} (1 - z_i) = \sum_{\emptyset \subsetneq I \subset \{1, \dots, r-1\}} \prod_{i \in I} (-1)^{|I|+1} (1 - z_i) -$$

$$\begin{aligned}
\sum_{I \subset \{1, \dots, r-1\}} \prod_{i \in I} (-1)^{|I|+1} (1-z_i)(1-z_r) &= 1 - \prod_{i=1}^{r-1} z_i + (1 - (1 - \prod_{i=1}^{r-1} z_i))(1-z_r) = \\
&= 1 - \prod_{i=1}^r z_i
\end{aligned}$$

□

A variation of this formula is the following:

Lemma 7.2. Set $t = \prod_{i=1}^h z_i \prod_{i=h+1}^r z_i^{-1}$, $a \in \mathbb{C}^*$

$$1-t = \sum_{\emptyset \subsetneq I \subset \{1, \dots, h\}} \prod_{i \in I} (-1)^{|I|+1} (1-z_i) - a^{-1} \sum_{\emptyset \subsetneq I \subset \{h+1, \dots, r\}} \prod_{i \in I} (-1)^{|I|+1} (1-z_i) +$$

$$(26) \quad a^{-1} \sum_{\emptyset \subsetneq I \subset \{h+1, \dots, r\}} \prod_{i \in I} (-1)^{|I|+1} (1-z_i)(1-at)$$

Proof. This is immediate from the previous Lemma once we remark that $1-t = 1 - \prod_{i=1}^h z_i - t(1 - \prod_{i=h+1}^r z_i)$. □

From this we get,

Lemma 7.3. Set $t = \prod_{i=1}^h z_i \prod_{i=h+1}^r z_i^{-1}$ with $0 \leq h \leq r$. Then

$$\begin{aligned}
\frac{1}{\prod_{i=1}^r (1-z_i)} &= \sum_{\emptyset \subsetneq I \subset \{1, \dots, h\}} \frac{(-1)^{|I|+1}}{(1-t) \prod_{i \notin I} (1-z_i)} - \\
(27) \quad &- \sum_{\emptyset \subsetneq I \subset \{h+1, \dots, r\}} \left(\frac{(-1)^{|I|+1}}{\prod_{i \notin I} (1-z_i)} - \frac{(-1)^{|I|+1}}{(1-t) \prod_{i \notin I} (1-z_i)} \right).
\end{aligned}$$

If $a \in \mathbb{C}^*$ and $a \neq 1$

$$\begin{aligned}
\frac{1}{\prod_{i=1}^r (1-z_i)(1-at)} &= \frac{a}{a-1} \left(\sum_{\emptyset \subsetneq I \subset \{1, \dots, h\}} \frac{(-1)^{|I|+1}}{(1-at) \prod_{i \notin I} (1-z_i)} - \right. \\
(28) \quad &- \frac{1}{a-1} \left(\sum_{\emptyset \subsetneq I \subset \{h+1, \dots, r\}} \left(\frac{(-1)^{|I|+1}}{\prod_{i \notin I} (1-z_i)} - \frac{(-1)^{|I|+1}}{(1-at) \prod_{i \notin I} (1-z_i)} \right) + \frac{1}{\prod_{i=1}^r (1-z_i)} \right).
\end{aligned}$$

Proof. The first relation follows from (26) dividing by $(1-t)(1-z_1)(1-z_2) \cdots (1-z_r)$ and taking $a = 1$.

The second writing

$$1 = \frac{a}{a-1}(1-t) - \frac{1}{a-1}(1-at)$$

and then dividing by $(1-at)(1-z_1)(1-z_2) \cdots (1-z_r)$. □

Let us now consider the group algebra $\mathbb{C}[\Gamma_{\mathbb{Q}}]$ of the vector space $\Gamma_{\mathbb{Q}}$, set $d = \prod_{(a, \chi) \in \Delta} (1 - a\chi)$ and $R = \mathbb{C}[\Gamma_{\mathbb{Q}}][1/d]$.

Lemma 7.4.

$$(29) \quad \frac{n}{1-x^n} = \sum_{i=0}^n \frac{1}{1-\zeta^i x}, \quad \zeta := e^{2\pi i/n}.$$

Proof. Take a variable t and derivatives with respect to t

$$\begin{aligned} \frac{nt^{n-1}}{t^n - x^n} dt &= d \log(t^n - x^n) = d \log\left(\prod_{i=0}^{n-1} (t - \zeta^i x)\right) \\ &= \sum_{i=0}^{n-1} d \log(t - \zeta^i x) = \sum_{i=0}^{n-1} \frac{1}{(t - \zeta^i x)} dt. \end{aligned}$$

Now set $t = 1$ in the coefficients of dt . □

Proposition 7.5. *Let*

$$\mathcal{S} := \{(b_1, \xi_1), \dots, (b_M, \xi_M)\}$$

be a sequence of elements compatible with Δ .

Let, for each $j = 1, \dots, M$, $\zeta_j \in \Gamma_{\mathbb{Q}}$ be such that there exists $0 \leq n_j \leq m_j$, $m_j > 0$, with $\zeta_j^{m_j} = \xi_j^{n_j}$ and set $\zeta = \zeta_1 \cdots \zeta_M$.

In R we can write the element

$$\frac{\zeta}{\prod_{i=1}^M (1 - b_i \xi_i)}$$

as a linear combination with constant coefficients of elements of the form

$$\frac{1}{(1 - c_1 \psi_1)^{h_1} \cdots (1 - c_r \psi_r)^{h_r}}$$

with $h_1, \dots, h_r \geq 0$ and $\{(c_1, \psi_1), \dots, (c_r, \psi_r)\}$ a non broken circuit.

Proof. Set $D_{\mathcal{S}} := \prod_{i=1}^M (1 - b_i \xi_i)$.

We first reduce to the case $\zeta = 1$. Indeed, for each j take $\theta_j \in \Gamma_{\mathbb{Q}}$ with $\theta_j^{m_j} = \xi_j$. We then have that $\zeta_j = \theta_j^{n_j}$. Also take a m_j -th root of b_j , f_j and write

$$(30) \quad 1 - b_j \xi_j = 1 - (f_j \theta_j)^{m_j} = \prod_{s=0}^{m_j-1} (1 - \exp(2\pi i s/m_j) f_j \theta_j)$$

Notice that for each s , the pair $(\exp(2\pi i s/m_j) f_j, \theta_j)$ is compatible with Δ . We now substitute in our sequence $\mathcal{S}$ the pair (b_j, ξ_j) with the sequence $\{(\exp(2\pi i s/m_j) f_j, \theta_j)\}$, $s = 0, \dots, m_j - 1$. We obtain a new sequence

$$\mathcal{S}' = \{(d_1, \mu_1), \dots, (d_N, \mu_N)\}$$

with $N = m_1 + m_2 + \cdots + m_M$ elements and with $D_{\mathcal{S}'} = D_{\mathcal{S}}$. In this equality, we have replaced (c_j, ξ_j) with a sequence of m_j pairs each having as second coordinate θ_j .

Since $\zeta_j = \theta_j^{n_j}$ and $n_j \leq m_j$, we have that $\zeta = \mu_1^{\varepsilon_1} \cdots \mu_N^{\varepsilon_N}$ with $\varepsilon_i \in \{0, 1\}$ for each i .

If $\zeta \neq 1$, let j_ζ be the least integer such that $\varepsilon_{j_\zeta} = 1$. Clearly either $\zeta' := \mu_{j_\zeta}^{-1}\zeta$ equals 1 or $j_{\zeta'} > j_\zeta$.

Adding and subtracting $(d_{j_\zeta}\mu_{j_\zeta})^{-1}\zeta$, a simple computation shows that

$$\frac{\zeta}{D_S} = -d_{j_\zeta}^{-1} \frac{\zeta'}{D_{S''}} + d_{j_\zeta}^{-1} \frac{\zeta'}{D_S}$$

where $S'' = S' - \{(d_{j_\zeta}, \mu_{j_\zeta})\}$. If $\zeta' = 1$ we are done. Otherwise, everything follows by induction on the number of indices j for which $\varepsilon_j = 0$.

Having reduced to the case $\zeta = 1$, let us now prove our claim for $1/D_S$. Set $\text{Supp}(\mathcal{S})$ equal to the subset in $\mathbb{C}^* \times \Gamma_{\mathbb{Q}}$ consisting of pairs (c, ξ) such that $(c, \xi) = (c_j, \xi_j)$ for some $j = 1, \dots, M$. If $\text{Supp}(\mathcal{S})$ is a non broken circuit there is nothing to prove.

Assume that the first condition in the definition of a no broken circuit is not verified i.e. there is a pair $(a, \chi) \in \Delta$ and distinct elements $(b_{i_1}, \xi_{i_1}) < \dots < (b_{i_t}, \xi_{i_t})$ in $\mathcal{S}$ with $(a, \chi) < (b_{i_1}, \xi_{i_1})$ and

$$(31) \quad (a\chi)^m (b_{i_1}\xi_{i_1})^{n_1} \dots (b_{i_t}\xi_{i_t})^{n_t} = 1$$

for suitable non zero integers $m, n_1, \dots, n_t$. In particular we get, since the elements χ and ξ_j are characters

$$(32) \quad \chi^m \xi_{i_1}^{n_1} \dots \xi_{i_t}^{n_t} = 1$$

Set $p = |mn_1 \dots n_t|$, and take $\theta_0, \theta_1, \dots, \theta_t$ in $\mathbb{C}[\Gamma_{\mathbb{Q}}]$ with the property that $\theta_0^{p/|m|} = \chi$, $\theta_h^{p/|n_h|} = \xi_{i_h}$ for each $h = 1, \dots, t$. Relation (32) becomes

$$(33) \quad (\theta_0^{\varepsilon_0} \dots \theta_t^{\varepsilon_t})^p = 1.$$

with $\varepsilon_j \in \{1, -1\}$. Hence

$$(34) \quad \theta_0^{\varepsilon_0} \dots \theta_t^{\varepsilon_t} = 1.$$

Apply, for each $1 \leq s \leq t$, the formula 7.4, for $x = c_s \theta_s$, $n = p/|n_s|$. Here c_s is a n -th root of b_{i_s} and substitute it in $1/D_S$. We get an expression of $1/D_S$ as a linear combination of p^t terms each of the form $1/D_{S'}$ where S' is obtained from $\mathcal{S}$ substituting each pair (b_{i_s}, ξ_{i_s}) with a pair (c_s, θ_s) , where as above c_s is a $p/|n_s|$ -th root of b_{i_s} . In particular S' has the same cardinality of $\mathcal{S}$ and the sequence of elements in Δ corresponding to the elements in S' and $\mathcal{S}$ coincide. Fix such a S' and take c_0 equal to the $p/|m|$ -th root of a such that

$$(35) \quad (c_0 \theta_0)^{\varepsilon_0} \dots (c_t \theta_t)^{\varepsilon_t} = 1.$$

We can then apply formula (27) of Lemma 7.3 and express $1/D_{S'}$ as a linear combination of elements of the form $1/D_{S''}$ where either the cardinality of S'' is strictly smaller than that of S' , or it is the same but S'' is obtained from S' removing a pair (c_s, θ_s) and inserting the smaller pair (c_0, θ_0) .

Assume that the second condition in the definition of a no broken circuit is not verified i.e. there are distinct elements $(b_{i_0}, \xi_{i_0}) < \dots < (b_{i_t}, \xi_{i_t})$ in $\mathcal{S}$

with $\xi_{i_1}, \dots, \xi_{i_t}$ linearly dependent. i.e.

$$(36) \quad \xi_{i_0}^{n_0} \cdots \xi_{i_t}^{n_t} = 1$$

for suitable integers $n_0, \dots, n_t$. Also we can assume that

$$(37) \quad b_{i_0}^{n_0} \cdots b_{i_t}^{n_t} \neq 1$$

otherwise we can use the previous discussion with $(a, \chi) = (b_{i_0}, \xi_{i_0})$. As before, set $p = |n_0 \dots n_t|$, and take $\theta_0, \dots, \theta_t$ in $\mathbb{C}[\Gamma_{\mathbb{Q}}]$ with the property that $\theta_h^{p/|n_h|} = \xi_{i_h}$ for each $h = 0, \dots, t$. Relation (36) implies

$$(38) \quad \theta_0^{\varepsilon_0} \cdots \theta_t^{\varepsilon_t} = 1.$$

Applying as before Lemma (7.4), for $x = c_s \theta_s$, $n = p/|n_s|$. Again c_s being a n -th root of b_{i_s} , for each $0 \leq s \leq t$. Substituting in $1/D_{\mathcal{S}}$, we get an expression of $1/D_{\mathcal{S}}$ as a linear combination of p^{t+1} terms each of the form $1/D_{\mathcal{S}'}$ where $\mathcal{S}'$ is obtained from $\mathcal{S}$ substituting each pair (b_{i_s}, ξ_{i_s}) with a pair (c_s, θ_s) , with c_s a $p/|n_s|$ -th root of b_{i_s} .

From relation (37) we deduce that

$$(39) \quad (c_0 \theta_0)^{\varepsilon_0} \cdots (c_t \theta_t)^{\varepsilon_t} = a \neq 1.$$

$a \in \mathbb{C}^*$. Now we can apply formula (28) in Lemma 7.3 (with $t = (c_0 \theta_0)^{-\varepsilon_0}$) and express $1/D_{\mathcal{S}'}$ as a linear combination of elements of the form $1/D_{\mathcal{S}''}$ where the cardinality of $\mathcal{S}''$ is strictly smaller than that of $\mathcal{S}'$. A simple induction then gives our Proposition. $\square$

7.2. The counting formula. We complete the argument following very closely the proof given in [26].

The setting is thus the following, we have a linear form ϕ with $\langle \phi, \alpha_i \rangle > 0$, $\forall i$. Setting $v_i := \frac{\alpha_i}{\langle \phi, \alpha_i \rangle}$, the vectors $\Psi := \{v_i\}$ span the r -dimensional real vector space V and lie in the affine hyperplane Π of equation $\langle \phi, x \rangle = 1$.

The intersection of the cone $C(\Psi) = C(\Delta)$ with Π is the convex polytope Σ envelop of the vectors v_i . Each cone, generated by $k+1$ independent vectors in Ψ (or of Δ), intersects Π in a k dimensional simplex.

We have a configuration of cones obtained by projecting a configuration of simplices and there is a simple dictionary to express properties of cones in terms of simplices and conversely.

It is well known that Σ is the union of the simplices with vertices independent vectors of Ψ . It is natural to define *regular* a point in Σ which is not contained in any $r-2$ dimensional simplex (or in the corresponding cone). The connected components of the set of regular points are called in [4] the *big cells*. We complete the set of cells with their boundary cells getting a stratification of Σ into cells and a stratification $\mathfrak{S}$ of $C(\Delta)$ into polyhedral cones.

Recall in fact that by [10], Theorem 5.5 we have .

Proposition 7.6. *Two elements of $C(\Delta)$ are in the same cone of $\mathfrak{S}$ if and only if they are contained in the same set of simplicial cones generated by no broken circuit bases.*

Remark The interesting feature of this statement is that, while $\mathfrak{S}$ is intrinsically defined, the no broken circuit bases depend on a choice of a total ordering of Δ .

We fix once and for all an orientation of the vector space $\Lambda \otimes_{\mathbb{Z}} \mathbb{Q}$ and take the invariant r -form ω_T defined in (12).

Definition 6. *Given a chamber $\mathfrak{c}$ we define the Jeffrey–Kirwan residue, relative to $\mathfrak{c}$, of a function $f \in R_T$ to be:*

$$JK(\mathfrak{c}, f) = (-1)^r \sum_P \sum_S \epsilon_S \text{res}_{S,P}(f\omega_T),$$

where P runs over the points of the arrangement and S on the no broken circuit bases in Δ_P such that $\mathfrak{c} \subset C(S)$ and ϵ_S equals 1, if S is equioriented with respect to our choice of orientation, -1 otherwise.

Remarks 7.7. 1) *Observe that $JK(\mathfrak{c}, f)$ does not depend on the choice of the orientation.*

2) *It follows from Proposition 4.7 that $JK(\mathfrak{c}, \chi^{-1}f)$ as function of χ is a periodic polynomial.*

Lemma 7.8. *Let $\pi : U \rightarrow T$ be a finite covering on T of degree m . Let $f \in R_T$ then $f \circ \pi \in R_U$, and*

$$(40) \quad JK(\mathfrak{c}, f) = JK(\mathfrak{c}, f \circ \pi)$$

Proof. A non broken circuit S associated to a point P in T is also associated to each point in $\pi^{-1}(P)$ in U . In this way one obtains all points in U associated to S . Also from the definition of the local residue and Lemma 4.10, we deduce that if $Q \in \pi^{-1}(P)$

$$\text{res}_{S,Q}(f\omega_U) = m^{-1} \text{res}_{S,P}(f\omega_T).$$

Since $m = |\pi^{-1}(P)|$ everything follows. $\square$

Lemma 7.9. 1) *Consider a non broken circuit basis $S := \{\psi_1, \dots, \psi_r\}$, a character $\mu = \prod_i \psi_i^{k_i}$ with $0 < k_i \in \mathbb{N}$ and a function*

$$\gamma := \frac{\mu^{-1}}{\prod_{i=1}^k (1 - a_{i_1} \psi_{i_k})^{h_i}}, \quad 1 \leq i_1 < \dots < i_k \leq r.$$

Then

$$(41) \quad \text{res}(\gamma) = (-1)^r \epsilon_S \sum_P \text{res}_{S,P}(\gamma\omega),$$

where P lies in the finite set of points associated to S (i.e. defined by the equations $1 - a_i \psi_i = 0$, $i = 1, \dots, r$). In particular if $k < r$ both terms are zero.

2) If μ is regular and $\mathfrak{c}$ is the unique chamber containing μ ,

$$(-1)^r \epsilon_S \sum_P \text{res}_{S,P}(\gamma\omega) = JK(\mathfrak{c}, \gamma).$$

The sum being on the set of P associated to S .

Proof. 1) Taking as Δ the set $\Delta_S = \{(a_1, \psi_1), \dots, (a_r, \psi_r)\}$ we see that

$$JK(C(S), \gamma) = (-1)^r \epsilon_S \sum_P \text{res}_{S,P}(\gamma\omega),$$

so by Lemma 7.8 the right hand side of formula (41) is independent of a torus on which the ψ_i are defined. By definition the same is true for $\text{res}(\gamma)$. Thus we can assume we are on the torus for which S is a basis of the character group. In this case, there is a unique point P associated to S . Furthermore $\omega = \epsilon_S d \log \psi_1 \wedge \dots \wedge d \log \psi_r$. Now one separates the variables and reduces to the 1 dimensional case. This is an elementary immediate instance of the knapsack problem (see §1.1 or [26]).

2) We need to see that, if (S', Q) is not one of our (S, P) , then $\text{res}_{S',Q}(\gamma\omega) = 0$. If $Q \neq P$ at least one on the factors $1 - a_i \psi_i$ is holomorphic in Q and we can apply part 1) of Theorem 4.5. If $P = Q$ remark that the form $\gamma\omega$ is in the image of the cohomology of $\mathcal{A}_{\Delta_S} \supset \mathcal{A}_\Delta$. Our claim now follows from formula (13) of Theorem 4.6 applied to $P \in \mathcal{A}_{\Delta_S}$. $\square$

Theorem 7.10. *Let*

$$f := \frac{1}{\prod_{(a,\chi) \in \Delta} (1 - a\chi)^{h_{a,\chi}}}, \quad h_{a,\chi} \in \mathbb{N}$$

If β is in the closure of a chamber $\mathfrak{c}$:

$$(42) \quad c_\beta = \text{res}(\beta^{-1} f \omega) = JK(\mathfrak{c}, \beta^{-1} f)$$

Proof. Using Lemma 7.8, we can pass to any finite covering U of T . For a suitable covering we can find a character ξ such that $\beta\xi$ lies in the interior of $\mathfrak{c}$ and the function ξf has an expansion as in Proposition 7.5. Having done this, we are reduced to prove our identity for each single term of this expansion. Write $\beta^{-1} f = (\beta\xi)^{-1} \xi f$. Each term of the expansion of ξf satisfies the hypotheses of the previous lemma with $\mu = \beta\xi$. The result follows. $\square$

Remarks: From the remark after Definition 7.2 follows that in the previous setting, c_β as function of β is a periodic polynomial. It has also a remarkable continuity property. That is if β is in the boundary of two different chambers there are two possibly different periodic polynomials on these chambers which agree on the intersection of their closure, in particular on β .

REFERENCES

- [1] W. Baldoni-Silva, J.A. De Loera, M. Vergne, *Counting Integer flows in Networks*, Found. Comput. Math. 4 (2004), no. 3, 277–314.

- [2] W. Baldoni-Silva, M. Vergne, *Residues formulae for volumes and Ehrhart polynomials of convex polytopes*, preprint math.CO/0103097
- [3] E.T. Bell, *Interpolated denumerants and Lambert series*, Amer. J. Math. 65 (1943), 382–386.
- [4] M. Brion, M. Vergne, *Residue formulae, vector partition functions and lattice points in rational polytopes*. J.A.M.S. (v. 10) 4 (1997), 797–833.
- [5] M. Brion, M. Vergne, *Arrangement of hyperplanes. I. Rational functions and Jeffrey-Kirwan residue*. Ann. Sci. cole Norm. Sup. (4) 32 (1999), no. 5, 715–741.
- [6] S. C. Coutinho, *A primer of algebraic D-modules*. London Mathematical Society Student Texts, 33. Cambridge University Press, Cambridge, 1995.
- [7] W. Dahmen, C. Micchelli, *The number of solutions to linear Diophantine equations and multivariate splines*. Trans. Amer. Math. Soc. 308 (1988), no. 2, 509–532.
- [8] C. DeConcini, C. Procesi, *Wonderful Models of subspace arrangements*. Selecta Math. (N.S.) 1 (1995), no. 3, 459–494.
- [9] C. DeConcini, C. Procesi, *Nested sets and Jeffrey Kirwan residues*. in: Geometric methods in Algebra and Number Theory, Bogomolov, F., Tschinkel. Y., Eds., vol. 235, 139–150.
- [10] C. DeConcini, C. Procesi, *On the geometry of graph arrangements*. math.CO/0412130
- [11] A. Dimca, G.I. Lehrer, *Purity and equivariant weight polynomials*. Algebraic groups and Lie groups, 161–181, Austral. Math. Soc. Lect. Ser., 9, Cambridge Univ. Press, Cambridge, 1997
- [12] J.M. Douglass, *Toral arrangements and hyperplane arrangements*. Rocky Mountain J. Math. 28 (1998), no. 3, 939–956.
- [13] E. Ehrhart, *Sur un problème de géométrie diophantienne linéaire I,II*, J. Reine Angew. Math., 1967, 226, pp. 1–29; 1967, 227, pp. 25–49.
- [14] R. Hartshorne, *Local cohomology*. A seminar given by A. Grothendieck, Harvard University, Fall, 1961. Lecture Notes in Mathematics, No. 41 Springer-Verlag, Berlin-New York 1967 vi+106 pp.
- [15] A. Henderson, *The symmetric group representation on cohomology of the regular elements of a maximal torus of the special linear group*, preprint <http://www.maths.usyd.edu.au/u/anthonyh/>
- [16] A. Grothendieck, *On the de Rham cohomology of algebraic varieties*. Inst. Hautes Études Sci. Publ. Math. No. 29 (1966), 95–103.
- [17] Lehrer, G. I. *The cohomology of the regular semisimple variety*. J. Algebra 199 (1998), no. 2, 666–689.
- [18] Lehrer, G. I. *A toral configuration space and regular semisimple conjugacy classes*. Math. Proc. Cambridge Philos. Soc. 118 (1995), no. 1, 105–113.

- [19] E. Looijenga, *Cohomology of $\mathcal{M}_3$ and $\mathcal{M}_3^1$* . Mapping class groups and moduli spaces of Riemann surfaces (Göttingen, 1991/Seattle, WA, 1991), Contemp. Math., 150, Amer. Math. Soc., 1993, 205–228.
- [20] C. Macmeikan, *Modules of derivations for toral arrangements*. Indag. Math. (N.S.) 15 (2004), no. 2, 257–267.
- [21] C. Macmeikan, *The Poincaré polynomial of an mp arrangement*. Proc. Amer. Math. Soc. 132 (2004), no. 6, 1575–1580.
- [22] R. Mac Pherson, C. Procesi, *Making conical compactifications into wonderful ones*. Selecta Math. 4, (1998), n.1, 125–139. Dekker 1998.
- [23] P. Orlik, H. Terao, *Arrangements of hyperplanes*. Grundlehren der Mathematischen Wissenschaften 300. Springer-Verlag, Berlin, 1992.
- [24] A. Szenes, *Iterated residues and multiple Bernoulli polynomials*. Int. Math. Res. Not. (1998), No.18, 937–956.
- [25] A. Szenes, *Residue theorem for rational trigonometric sums and Verlinde’s formula*. Duke Math. J. 118 (2003), no. 2, 189–227.
- [26] A. Szenes, M. Vergne, *Residue formulae for vector partitions and Euler-Maclaurin sums* Formal power series and algebraic combinatorics (Scottsdale, AZ, 2001). Adv. in Appl. Math. 30 (2003), no. 1-2, 295–342.
- [27] A. Szenes, M. Vergne, *Toric reduction and a conjecture of Batyrev and Materov* Invent. Math. 158 (2004), no. 3, 453–495.

DIP. MAT. CASTELNUOVO, UNIV. DI ROMA LA SAPIENZA, ROME, ITALY
E-mail address: `deconcini@mat.uniroma1.it`

DIP. MAT. CASTELNUOVO, UNIV. DI ROMA LA SAPIENZA, ROME, ITALY
E-mail address: `procesi@mat.uniroma1.it`